

Trying 31060000009999...Open

DIALOG INFORMATION SERVICES

PLEASE LOGON:

\*\*\*\*\* HHHHHHHH SSSSSSSS? ### Status: Signing onto Dialog \*\*\*\*\*

ENTER PASSWORD:

\*\*\*\*\* HHHHHHHH SSSSSSSS? \*\*\*\*\*

Welcome to DIALOG

### Status: Login successfulDialog level 05.15.00D

Last logoff: 05jan07 15:02:16

Logon file405 23jan07 12:45:47

\*\*\* ANNOUNCEMENTS \*\*\*

\*\*\*

NEW FILES RELEASED

\*\*\*Engineering Index Backfile (File 988)

\*\*\*Verdict Market Research (File 769)

\*\*\*EMCare (File 45)

\*\*\*Trademarkscan - South Korea (File 655)

RESUMED UPDATING

\*\*\*File 141, Reader's Guide Abstracts

\*\*\*

RELOADS COMPLETED

\*\*\*Files 340, 341 & 942, CLAIMS/U.S. Patents - 2006 reload now online

\*\*\*Files 173 & 973, Adis Clinical Trials Insight

\*\*\*File 11, PsycInfo

\*\*\*File 531, American Business Directory

\*\*\*

DATABASES REMOVED

\*\*\*File 196, FINDEX

\*\*\*File 468, Public Opinion Online (POLL)

Chemical Structure Searching now available in Prous Science Drug

Data Report (F452), Prous Science Drugs of the Future (F453),

IMS R&D Focus (F445/955), Pharmaprojects (F128/928), Beilstein

Facts (F390), Derwent Chemistry Resource (F355) and Index Chemicus

(File 302).

\*\*\*

>>>For the latest news about Dialog products, services, content<<<

>>>and events, please visit What's New from Dialog at <<<

>>><http://www.dialog.com/whatsnew/>. You can find news about<<<

>>>a specific database by entering HELP NEWS <file number>.<<<

\* \* \*

SYSTEM:HOME

Cost is in DialUnits

Menu System II: D2 version 1.8.0 term=ASCII

\*\*\* DIALOG HOMEBASE(SM) Main Menu \*\*\*

Information:

1. Announcements (new files, reloads, etc.)
2. Database, Rates, & Command Descriptions
3. Help in Choosing Databases for Your Topic
4. Customer Services (telephone assistance, training, seminars, etc.)
5. Product Descriptions

Connections:

6. DIALOG(R) Document Delivery
7. Data Star(R)

(c) 2003 Dialog, a Thomson business. All rights reserved.

/H = Help

/L = Logoff

/NOMENU = Command Mode

Enter an option number to view information or to connect to an online service. Enter a BEGIN command plus a file number to search a database (e.g., B1 for ERIC).

?

Terminal set to DLINK

\*\*\* DIALOG HOMEBASE(SM) Main Menu \*\*\*

Information:

1. Announcements (new files, reloads, etc.)
2. Database, Rates, & Command Descriptions
3. Help in Choosing Databases for Your Topic
4. Customer Services (telephone assistance, training, seminars, etc.)
5. Product Descriptions

Connections:

6. DIALOG(R) Document Delivery
7. Data Star(R)

(c) 2003 Dialog, a Thomson business. All rights reserved.

/H = Help

/L = Logoff

/NOMENU = Command Mode

Enter an option number to view information or to connect to an online service. Enter a BEGIN command plus a file number to search a database (e.g., B1 for ERIC).

? b biosci

```
>>>          44 is unauthorized
>>>          76 is unauthorized
>>>2 of the specified files are not available
    23jan07 12:45:52 User276653 Session D80.1
        $0.00    0.245 DialUnits FileHomeBase
    $0.00 Estimated cost FileHomeBase
    $0.02 TELNET
    $0.02 Estimated cost this search
    $0.02 Estimated total session cost    0.245 DialUnits
```

SYSTEM:OS - DIALOG OneSearch

- File 5:Biosis Previews(R) 1969-2007/Jan W2  
(c) 2007 The Thomson Corporation
- File 24:CSA Life Sciences Abstracts 1966-2007/Nov  
(c) 2007 CSA.
- File 28:Oceanic Abstracts 1966-2007/Nov  
(c) 2007 CSA.
- File 34:SciSearch(R) Cited Ref Sci 1990-2007/Jan W2  
(c) 2007 The Thomson Corp
- File 35:Dissertation Abs Online 1861-2006/Nov

(c) 2006 ProQuest Info&Learning  
File 40:Enviroline(R) 1975-2007/Dec  
(c) 2007 Congressional Information Service  
File 41:Pollution Abstracts 1966-2007/Nov  
(c) 2007 CSA.  
File 45:EMCare 2007/Jan W2  
(c) 2007 Elsevier B.V.  
File 50:CAB Abstracts 1972-2007/Dec  
(c) 2007 CAB International  
File 65:Inside Conferences 1993-2007/Jan 23  
(c) 2007 BLDSC all rts. reserv.  
File 71:ELSEVIER BIOBASE 1994-2007/Jan W3  
(c) 2007 Elsevier B.V.  
File 73:EMBASE 1974-2007/Jan 17  
(c) 2007 Elsevier B.V.  
**\*File 73: Elsevier will not provide the daily update to Embase on January 18. Tomorrow's update will contain both days.**  
File 91:MANTIS(TM) 1880-2006/Jan  
2001 (c) Action Potential  
File 94:JICST-EPlus 1985-2007/Jan W3  
(c)2007 Japan Science and Tech Corp(JST)  
**\*File 94: UD200609W2 is the last update for 2006. UD200701W1 is the first update for 2007. The file is complete and up to date.**  
File 98:General Sci Abs 1984-2007/Jan  
(c) 2007 The HW Wilson Co.  
File 110:WasteInfo 1974-2002/Jul  
(c) 2002 AEA Techn Env.  
**\*File 110: This file is closed (no updates)**  
File 135:NewsRx Weekly Reports 1995-2007/Jan W2  
(c) 2007 NewsRx  
File 136:BioEngineering Abstracts 1966-2007/Nov  
(c) 2007 CSA.  
File 143:Biol. & Agric. Index 1983-2007/Dec  
(c) 2007 The HW Wilson Co  
File 144:Pascal 1973-2007/Jan W2  
(c) 2007 INIST/CNRS  
File 155:MEDLINE(R) 1950-2006/Dec 16  
(c) format only 2006 Dialog  
**\*File 155: MEDLINE has resumed updating with UD20061209. Please see HELP NEWS 154 for details.**  
File 164:Allied & Complementary Medicine 1984-2007/Jan  
(c) 2007 BLHCIS  
File 172:EMBASE Alert 2007/Jan 23  
(c) 2007 Elsevier B.V.  
File 185:Zoological Record Online(R) 1978-2007/Feb  
(c) 2007 The Thomson Corp.  
File 357:Derwent Biotech Res. \_1982-2007/Jan W1  
(c) 2007 The Thomson Corp.  
File 369:New Scientist 1994-2007/Oct W3  
(c) 2007 Reed Business Information Ltd.  
File 370:Science 1996-1999/Jul W3  
(c) 1999 AAAS  
**\*File 370: This file is closed (no updates). Use File 47 for more current information.**  
File 391:Beilstein Reactions 2006/Q4  
(c) 2006 Beilstein GmbH  
File 434:SciSearch(R) Cited Ref Sci 1974-1989/Dec

(c) 2006 The Thomson Corp  
File 467:ExtraMED(tm) 2000/Dec  
(c) 2001 Informania Ltd.

Set	Items	Description
---	-----	-----
? s	crystallin	
	S1 40380	CRYSTALLIN
? s	arginine(n)chloride	
	416125	ARGININE
	1978652	CHLORIDE
	S2 483	ARGININE (N) CHLORIDE
? s	s1 and s2	
	40380	S1
	483	S2
	S3 1	S1 AND S2
? t	s3/9,k/all	

3/9,K/1 (Item 1 from file: 73)  
DIALOG(R)File 73:EMBASE  
(c) 2007 Elsevier B.V. All rts. reserv.

13167229 EMBASE No: 2005234088  
**Alexander-disease mutation of GFAP causes filament disorganization and decreased solubility of GFAP**  
Hsiao V.C.; Tian R.; Long H.; Perng M.D.; Brenner M.; Quinlan R.A.; Goldman J.E.  
J.E. Goldman, Department of Pathology, Center for Neurobiology and Behavior, Columbia University, New York, NY 10032 United States  
AUTHOR EMAIL: jeg5@columbia.edu  
Journal of Cell Science ( J. CELL SCI. ) (United Kingdom) 01 MAY 2005, 118/9 (2057-2065)  
CODEN: JNCSA ISSN: 0021-9533  
DOCUMENT TYPE: Journal ; Article  
LANGUAGE: ENGLISH SUMMARY LANGUAGE: ENGLISH  
NUMBER OF REFERENCES: 66

Alexander disease is a fatal neurological illness characterized by white-matter degeneration and the formation of astrocytic cytoplasmic inclusions called Rosenthal fibers, which contain the intermediate filament glial fibrillary acidic protein (GFAP), the small heat-shock proteins HSP27 and alphaB- crystallin , and ubiquitin. Many Alexander-disease patients are heterozygous for one of a set of point mutations in the GFAP gene, all of which result in amino acid substitutions. The biological effects of the most common alteration, R239C, were tested by expressing the mutated protein in cultured cells by transient transfection. In primary rat astrocytes and Cos-7 cells, the mutant GFAP was incorporated into filament networks along with the endogenous GFAP and vimentin, respectively. In SW13VimSUP- cells, which have no endogenous cytoplasmic intermediate filaments, wild-type human GFAP frequently formed filamentous bundles, whereas the R239C GFAP formed 'diffuse' and irregular patterns. Filamentous bundles of R239C GFAP were sometimes formed in SW13VimSUP- cells when wild-type GFAP was co-transfected. Although the presence of a suitable coassembly partner (vimentin or GFAP) reduced the potential negative effects of the R239C mutation on GFAP network formation, the mutation affected the stability of GFAP in cells in a dominant fashion. Extraction of transfected SW13VimSUP- cells with Triton-X-100-containing buffers

showed that the mutant GFAP was more resistant to solubilization at elevated KCl concentrations. Both wild-type and R239C GFAP assembled into 10 nm filaments with similar morphology in vitro. Thus, although the R239C mutation does not appear to affect filament formation per se, the mutation alters the normal solubility and organization of GFAP networks.

#### DRUG DESCRIPTORS:

\*glial fibrillary acidic protein--endogenous compound--ec  
heat shock protein 27--endogenous compound--ec; alpha **crystallin**  
--endogenous compound--ec; beta **crystallin** --endogenous compound--ec;  
ubiquitin--endogenous compound--ec; amino acid--endogenous compound--ec;  
mutant protein--endogenous compound--ec; vimentin--endogenous compound--ec;  
triton x 100; buffer; potassium **chloride** ; **arginine** --endogenous compound  
--ec; cysteine--endogenous compound--ec

#### MEDICAL DESCRIPTORS:

\*Alexander disease--diagnosis--di; \*Alexander disease--etiology--et  
gene mutation; neurologic disease--etiology--et; clinical feature; white  
matter; astrocyte; cell inclusion; heterozygosity; point mutation; amino  
acid substitution; protein expression; cell culture; genetic transfection;  
cytoplasm; wild type; protein assembly; intermediate filament; protein  
stability; solubilization; concentration (parameters); cell structure; in  
vitro study; protein structure; human; nonhuman; rat; controlled study;  
human cell; animal cell; article; priority journal

CAS REGISTRY NO.: 60267-61-0 (ubiquitin); 65072-01-7 (amino acid);  
7447-40-7 (potassium chloride); 1119-34-2, 15595-35-4, 7004-12-8,  
74-79-3 (arginine); 4371-52-2, 52-89-1, 52-90-4 (cysteine)

#### SECTION HEADINGS:

005 General Pathology and Pathological Anatomy  
008 Neurology and Neurosurgery  
022 Human Genetics  
029 Clinical and Experimental Biochemistry

...intermediate filament glial fibrillary acidic protein (GFAP), the  
small heat-shock proteins HSP27 and alphaB- **crystallin** , and ubiquitin.  
Many Alexander-disease patients are heterozygous for one of a set of point

...

#### DRUG DESCRIPTORS:

heat shock protein 27--endogenous compound--ec; alpha **crystallin**  
--endogenous compound--ec; beta **crystallin** --endogenous compound--ec;  
ubiquitin--endogenous compound--ec; amino acid--endogenous compound--ec;  
mutant protein--endogenous compound--ec; vimentin--endogenous compound--ec;  
triton x 100; buffer; potassium **chloride** ; **arginine** --endogenous compound  
--ec; cysteine--endogenous compound--ec

? s arginine and chloride

416125 ARGININE

1978652 CHLORIDE

S4 12788 ARGININE AND CHLORIDE

? s s1 and s4

40380 S1

12788 S4

S5 10 S1 AND S4

? t s5/9,k/1-10

5/9,K/1 (Item 1 from file: 34)

DIALOG(R) File 34:SciSearch(R) Cited Ref Sci

(c) 2007 The Thomson Corp. All rts. reserv.

05956303 Genuine Article#: XK290 Number of References: 42

**Title: Expression of arginine decarboxylase in seedlings of indica rice (*Oryza sativa* L) cultivars as affected by salinity stress**

Author(s): Chattopadhyay MK; Gupta S; Sengupta DN; Ghosh B (REPRINT)

Corporate Source: BOSE INST,DEPT BOT, 93-1 APC RD/CALCUTTA 700009/W  
BENGAL/INDIA/ (REPRINT); BOSE INST,DEPT BOT/CALCUTTA 700009/W  
BENGAL/INDIA/

Journal: PLANT MOLECULAR BIOLOGY, 1997, V34, N3 (JUN), P477-483

ISSN: 0167-4412 Publication date: 19970600

Publisher: KLUWER ACADEMIC PUBL, SPUIBOULEVARD 50, PO BOX 17, 3300 AA  
DORDRECHT, NETHERLANDS

Language: English Document Type: ARTICLE

Geographic Location: INDIA

Subfile: CC LIFE--Current Contents, Life Sciences; CC AGRI--Current  
Contents, Agriculture, Biology & Environmental Sciences

Journal Subject Category: PLANT SCIENCES; BIOCHEMISTRY & MOLECULAR BIOLOGY

**Abstract:** The effect of salinity stress on the activity of **arginine** decarboxylase (ADC, EC 4.1.1.19), the first enzyme in biosynthesis of polyamines (PA) from **arginine**, as well as its transcript level has been compared in salt-sensitive (M-1-48) and salt-tolerant (Pokkali) rice cultivars. Treatment of 72 h grown seedlings either with increasing concentrations of NaCl or with 150 mM NaCl for different time periods, showed a gradual increase of activity in Pokkali. In M-1-48 an immediate increase followed by sharp decrease was observed on prolonged treatment beyond 6 h or above 150 mM NaCl. To generate a DNA probe for ADC, the polymerase chain reaction was used with oat genomic DNA and sequence-specific primers. A region of oat genomic DNA containing a coding sequence for 166 amino acids of the C-terminal part of the ADC enzyme was amplified and called OAD1. Southern analysis of EcoRI- or BamHI-cut genomic DNAs from different cultivars of rice with OAD1 as the probe revealed strong hybridization with one DNA fragment of rice and restriction fragment length polymorphism (RFLP) was noticed. Northern analysis of total RNA of rice with OAD1 as the probe revealed hybridization with a transcript of similar size to the ADC transcript in oat. While in Pokkali, at least a 20-fold accumulation of OAD1 homologous transcript was detected after treatment with 200 mM NaCl, only a seven-fold increase in transcript level was found in M-1-48 after 150 mM NaCl treatment. Results suggest that in the salt-tolerant rice cultivar Pokkali ADC enzyme activity increases and its transcript also accumulates during the prolonged salinity stress, this mechanism is absent in the salt-sensitive rice cultivar M-1-48 where a prolonged period of salinity stress down-regulates both ADC activity and its transcript level.

**Descriptors--Author Keywords:** **arginine** decarboxylase ; gene expression ; *Oryza sativa* ; polyamines ; rice ; salinity stress

**Identifiers--Key Word Plus(R):** POLYAMINE ACCUMULATION; OSMOTIC-STRESS; SALT TOLERANCE; WATER-STRESS; LEAVES; PUTRESCINE; RESPONSES; PLANTS; ACID; **CHLORIDE**

**Research Fronts:** 95-1488 001 (ORNITHINE DECARBOXYLASE; SPERMIDINE  
TRANSPORT IN HUMAN BREAST-CANCER CELLS; REGULATION OF CELLULAR  
POLYAMINES)

95-3190 001 (INCREASED ABUNDANCE OF SPECIFIC SKELETAL-MUSCLE  
PROTEIN-TYROSINE PHOSPHATASES; ALPHA-B-CRYSTALLIN EXPRESSION)

95-3260 001 (ABSCISIC-ACID RESPONSE ELEMENTS; STRESS PROTEINS; GENE IN  
ARABIDOPSIS-THALIANA; DIFFERENTIAL EXPRESSION; POTENTIAL REGULATION;  
DESICCATION TOLERANCE)

95-5061 001 (STRUCTURAL GENE; GLTC-DEPENDENT REGULATION OF

BACILLUS-SUBTILIS GLUTAMATE SYNTHASE EXPRESSION; ARABIDOPSIS TYPE-1  
PROTEIN PHOSPHATASE)

95-5565 001 (POLYAMINE BIOSYNTHESIS; DEVELOPMENT OF ZYGOTIC EMBRYOS;  
MOUSE ORNITHINE DECARBOXYLASE CDNA IN CARROT PROMOTES SOMATIC  
EMBRYOGENESIS)

Cited References:

AKBAR M, 1972, V22, P277, JAPANESE J BREEDING  
BASU R, 1988, V15, P777, AUST J PLANT PHYSIOL  
BASU R, 1991, V82, P575, PHYSIOL PLANTARUM  
BELL E, 1990, V224, P431, MOL GEN GENET  
BESFORD RT, 1993, V189, P201, PLANTA  
BORRELL A, 1996, V98, P105, PHYSIOL PLANTARUM  
BRADFORD MM, 1976, V72, P248, ANAL BIOCHEM  
BRAY EA, 1993, V103, P1035, PLANT PHYSIOL  
BROECK DV, 1994, V106, P559, PLANT PHYSIOL  
CHOMCZYNSKI P, 1987, V162, P156, ANAL BIOCHEM  
COLEMAN RG, 1956, V20, P393, ANN BOT-LONDON  
DAS S, 1987, V14, P108, PLANT PHYSIOL BIOCH  
FLORES H, 1982, V217, P1259, SCIENCE  
FLORES HE, 1984, V75, P102, PLANT PHYSIOL  
FLORES HE, 1990, V12, P217, STRESS RESPONSES PLA  
FRIEDMAN R, 1989, V76, P295, PHYSIOL PLANTARUM  
KALLIO A, 1981, V20, P3163, BIOCHEMISTRY-US  
KANDPAL RP, 1985, V11, P365, BIOCHEM INT  
KRISHNAMURTHY R, 1989, V91, P500, PLANT PHYSIOL  
KUEHN GD, 1990, P190, POLYAMINES ETHYLENE  
LEVITT J, 1980, P386, SALT STRESSES RESPON  
MALMBERG RL, 1994, V269, P2703, J BIOL CHEM  
MALMBERG RL, 1992, V100, P146, PLANT PHYSIOL  
MOONS A, 1995, V107, P177, PLANT PHYSIOL  
PRIEBE A, 1978, V12, P365, PLANT SCI LETT  
RAMAKRISHNA S, 1975, V59, P377, EUR J BIOCHEM  
RASTOGI R, 1993, V103, P829, PLANT PHYSIOL  
REGGIANI R, 1990, V31, P489, PLANT CELL PHYSIOL  
SAMBROOK J, 1989, MOL CLONING LAB MANU  
SHEVYAKOVA NI, 1985, V3, P365, PLANT GROWTH REGUL  
SHEVYAKOVA NI, 1981, V28, P594, SOV PLANT PHYSIOL  
SINDHU RK, 1979, V18, P1937, PHYTOCHEMISTRY  
SMITH TA, 1985, V36, P117, ANNU REV PLANT PHYS  
SMITH TA, 1973, V12, P2093, PHYTOCHEMISTRY  
SMITH TA, 1979, V18, P1447, PHYTOCHEMISTRY  
TABOR CW, 1984, V53, P749, ANNU REV BIOCHEM  
TIBURCIO AF, 1994, V22, P5455, BIOCHEM SOC T  
TIBURCIO AF, 1994, V45, P1789, J EXP BOT  
TURNER LB, 1986, V37, P170, J EXP BOT  
YANAGISAWA H, 1982, V21, P2201, PHYTOCHEMISTRY  
YOUNG ND, 1983, V71, P767, PLANT PHYSIOL  
ZHU H, 1993, V21, P5279, NUCLEIC ACIDS RES

Title: Expression of arginine decarboxylase in seedlings of indica rice  
(Oryza sativa L) cultivars as affected by salinity stress

Abstract: The effect of salinity stress on the activity of arginine  
decarboxylase (ADC, EC 4.1.1.19), the first enzyme in biosynthesis of  
polyamines (PA) from arginine, as well as its transcript level has  
been compared in salt-sensitive (M-1-48...

...Identifiers--POLYAMINE ACCUMULATION; OSMOTIC-STRESS; SALT TOLERANCE;  
WATER-STRESS; LEAVES; PUTRESCINE; RESPONSES; PLANTS; ACID; CHLORIDE

...Research Fronts: POLYAMINES)

95-3190 001 (INCREASED ABUNDANCE OF SPECIFIC SKELETAL-MUSCLE

PROTEIN-TYROSINE PHOSPHATASES; ALPHA-B- CRYSTALLIN EXPRESSION)

95-3260 001 (ABSCISIC-ACID RESPONSE ELEMENTS; STRESS PROTEINS; GENE IN ARABIDOPSIS-THALIANA; DIFFERENTIAL...

5/9,K/2 (Item 2 from file: 34)

DIALOG(R)File 34:SciSearch(R) Cited Ref Sci

(c) 2007 The Thomson Corp. All rts. reserv.

05799266 Genuine Article#: WY280 Number of References: 40

Title: Chronic ethanol increases N-methyl-D-aspartate-stimulated nitric oxide formation but not receptor density in cultured cortical neurons

Author(s): Chandler LJ (REPRINT) ; Sutton G; Norwood D; Sumners C; Crews FT

Corporate Source: LOUISIANA STATE UNIV,MED CTR, DEPT PHARMACOL, 1501 KINGS HIGHWAY/SHREVEPORT//LA/71130 (REPRINT); UNIV FLORIDA,COLL MED, DEPT PHYSIOL/GAINESVILLE//FL/32610; UNIV N CAROLINA,BOWLES CTR ALCOHOL STUDIES/CHAPEL HILL//NC/27599

Journal: MOLECULAR PHARMACOLOGY, 1997, V51, N5 (MAY), P733-740

ISSN: 0026-895X Publication date: 19970500

Publisher: WILLIAMS & WILKINS, 351 WEST CAMDEN ST, BALTIMORE, MD 21201-2436

Language: English Document Type: ARTICLE

Geographic Location: USA

Subfile: CC LIFE--Current Contents, Life Sciences

Journal Subject Category: PHARMACOLOGY & PHARMACY; BIOCHEMISTRY & MOLECULAR BIOLOGY

Abstract: The effects of prolonged ethanol exposure on excitatory amino acid receptor stimulated nitric oxide (NO) formation were examined in primary rat cortical neuronal cultures. Chronic ethanol (4 days, 100 mM) potentiated N-methyl-D-aspartate (NMDA)-stimulated NO formation as determined by measuring the conversion of [H-3] arginine to [H-3]citrulline. In contrast, chronic ethanol had no effect on NO formation stimulated by kainate, alpha-amino-3-hydroxy-5-methyl-4-isoxalonepropionic acid, or the calcium ionophore ionomycin. Potassium chloride -stimulated NO formation was also enhanced by chronic ethanol treatment, but this effect was not seen in the presence of the ionotropic glutamate receptor antagonists MK-801 and 6-cyano-7-nitroquinoxaline-2,3-dione. Immunoblot analysis of expression of NR1, NR2A, and NR2B receptor subunits showed no difference between control and chronic ethanol-treated cultures. In support of this apparent lack of change in receptor density, there was no difference in the specific binding of I-125-MK-801 between control and chronic ethanol-treated groups. These results demonstrate that prolonged ethanol exposure selectively enhanced NMDA receptor-stimulated NO formation, which may play an important role in alcohol dependence, withdrawal, and alcohol-associated brain damage. These results also suggest that chronic ethanol-induced increases in NMDA receptor function may not be due to a simple increase in the number of NMDA receptors or change in NMDA receptor subunit composition but may instead reflect more complicated and subtle changes.

Identifiers--KeyWord Plus(R): WITHDRAWAL SEIZURES; RAT-BRAIN; ALCOHOL-WITHDRAWAL; IONOPHORE COMPLEX; CHRONIC EXPOSURE; RAPID TOLERANCE; NMDA RECEPTORS; SYNTHASE; BINDING; SUBUNIT

Research Fronts: 95-3190 001 (INCREASED ABUNDANCE OF SPECIFIC



SKELETAL-MUSCLE PROTEIN-TYROSINE PHOSPHATASES; ALPHA-B- CRYSTALLIN  
EXPRESSION)

95-6313 001 (ETHANOL WITHDRAWAL; N-METHYL-D-ASPARTATE RECEPTORS; RAT  
HIPPOCAMPAL 2-DEOXYGLUCOSE UPTAKE IN-VITRO)

Cited References:

ADAMS ML, 1993, V17, P660, ALCOHOL CLIN EXP RES  
ADAMS ML, 1995, V19, P195, ALCOHOL CLIN EXP RES  
AHERN KV, 1994, V165, P211, NEUROSCI LETT  
CARTER LA, 1995, V64, P213, J NEUROCHEM  
CHANDLER LJ, 1993, V60, P1578, J NEUROCHEM  
CHANDLER LJ, 1994, V271, P67, J PHARMACOL EXP THER  
CHARNESS ME, 1983, V222, P1246, SCIENCE  
CHEN L, 1992, V356, P521, NATURE  
CHOU YC, 1990, V2, P29, J NEUROENDOCRINOL  
COLLINGRIDGE GL, 1989, V41, P143, PHARMACOL REV  
CREWS F, 1996, V39, P283, INT REV NEUROBIOL  
CREWS FT, 1993, P355, NIH933549 PUBL  
DAWSON TM, 1994, V14, P5247, J NEUROSCI  
FITZGERALD LW, 1995, V21, P93, SYNAPSE  
FOLLESA P, 1995, V29, P99, MOL BRAIN RES  
GRANT KA, 1990, V176, P289, EUR J PHARMACOL  
GULYA K, 1991, V547, P129, BRAIN RES  
HU XJ, 1995, V30, P347, MOL BRAIN RES  
IORIO KR, 1992, V41, P1142, MOL PHARMACOL  
KHANNA JM, 1993, V32, P43, BRAIN RES BULL  
KHANNA JM, 1995, V37, P599, BRAIN RES BULL  
KOLTCHINE V, 1993, V152, P13, NEUROSCI LETT  
KUMAR KN, 1991, V354, P70, NATURE  
LAEMMLI UK, 1970, V227, P680, NATURE  
LIEBERMAN DN, 1994, V369, P235, NATURE  
LILJEQUIST S, 1991, V192, P197, EUR J PHARMACOL  
MASOOD K, 1994, V45, P324, MOL PHARMACOL  
MONCADA S, 1993, V329, P2002, NEW ENGL J MED  
MORRISETT RA, 1990, V176, P103, EUR J PHARMACOL  
NAKANISHI S, 1992, V257, P597, SCIENCE WASHINGTON D  
ORTIZ J, 1995, V21, P289, SYNAPSE  
SEEBURG PH, 1993, V14, P297, TRENDS PHARMACOL SCI  
SNELL LD, 1993, V602, P91, BRAIN RES  
TREMWEI MF, 1994, V18, P1004, ALCOHOL CLIN EXP RES  
TREVISAN L, 1994, V62, P1635, J NEUROCHEM  
WANG LY, 1994, V369, P230, NATURE  
WANG YH, 1995, V65, P176, J NEUROCHEM  
WANG YT, 1994, V369, P233, NATURE  
WANG YT, 1996, V93, P1721, P NATL ACAD SCI USA  
WILLIAMS K, 1992, V42, P147, MOL PHARMACOL

...Abstract: D-aspartate (NMDA)-stimulated NO formation as determined by  
measuring the conversion of [H-3] **arginine** to [H-3]citrulline. In  
contrast, chronic ethanol had no effect on NO formation stimulated...

...alpha-amino-3-hydroxy-5-methyl-4-isoxalonepropionic acid, or the calcium  
ionophore ionomycin. Potassium **chloride** stimulated NO formation was  
also enhanced by chronic ethanol treatment, but this effect was not...

Research Fronts: 95-3190 001 (INCREASED ABUNDANCE OF SPECIFIC  
SKELETAL-MUSCLE PROTEIN-TYROSINE PHOSPHATASES; ALPHA-B- CRYSTALLIN  
EXPRESSION)

95-6313 001 (ETHANOL WITHDRAWAL; N-METHYL-D-ASPARTATE RECEPTORS; RAT

HIPPOCAMPAL 2-DEOXYGLUCOSE...

5/9,K/3 (Item 3 from file: 34)

DIALOG(R) File 34:SciSearch(R) Cited Ref Sci

(c) 2007 The Thomson Corp. All rts. reserv.

05765539 Genuine Article#: WW460 Number of References: 37

**Title: Simulatory effect of porcine insulin on noradrenaline secretion in guinea-pig ileum myenteric nerve terminals**

Author(s): Cheng JT (REPRINT) ; Hung CR; Lin MI

Corporate Source: NATL CHENG KUNG UNIV, COLL MED, DEPT PHARMACOL/TAINAN 70101//TAIWAN/ (REPRINT)

Journal: BRITISH JOURNAL OF PHARMACOLOGY, 1997, V121, N1 (MAY), P15-20

ISSN: 0007-1188 Publication date: 19970500

Publisher: STOCKTON PRESS, HOUNDMILLS, BASINGSTOKE, HAMPSHIRE, ENGLAND RG21 6XS

Language: English Document Type: ARTICLE

Geographic Location: TAIWAN

Subfile: CC LIFE--Current Contents, Life Sciences

Journal Subject Category: PHARMACOLOGY & PHARMACY; BIOCHEMISTRY & MOLECULAR BIOLOGY

**Abstract:** 1 The effect of insulin on the release of noradrenaline (NA) from nerve terminals was investigated in isolated ileal synaptosomes of guinea-pig. Release was determined as the amount of NA, quantified by h.p.l.c.-electrochemical detection, from samples incubated with insulin minus that in parallel blanks treated with some volume of vehicle.

2 Porcine insulin stimulated the secretion of NA in a concentration-dependent manner from 0.01 i.u. ml<sup>-1</sup>, while the value of lactate dehydrogenase in the incubated medium was not influenced by insulin.

3 The presence of insulin receptors in this preparation was illustrated by immunoblotting with insulin receptor monoclonal antibodies.

4 The release of NA by insulin was reduced by guanethidine and bretylium and it was markedly lowered in the samples obtained from guinea-pigs that had received an intraperitoneal injection of DSP-4, the noradrenergic neurotoxin.

5 Tetrodotoxin attenuated the action of insulin at concentrations sufficient to block sodium channels. The depolarizing effect of insulin on the membrane potential was also illustrated by a concentration-dependent increase in the fluorescence of bisoxonol, a potential-sensitive dye.

6 The action of insulin was attenuated by removal of calcium chloride from the bathing medium. The induction of calcium ion influx by insulin into the synaptosomes is supported by the inhibitory effects of the calcium channel blockers omega-conotoxin GVIA (for the N-type channels) and nifedipine (for the L-type channels).

7 These findings suggest that insulin can stimulate NA release from noradrenergic terminals via activation of calcium influx.

Descriptors--Author Keywords: insulin ; noradrenaline release ; bisoxonol ; calcium influx ; synaptosomal preparation of guinea-pig ileum

Identifiers--Keyword Plus(R): CYTOSOLIC CA-2+; HYPERTENSION; RATS;  
HYPERINSULINEMIA; CATECHOLAMINES; SYNAPTOSOMES; TETRODOTOXIN;  
ACTIVATION; INHIBITION; MEMBRANES

Research Fronts: 95-0917 002 (INSULIN-RESISTANCE IN SYSTEMIC  
HYPERTENSION; COMPENSATORY HYPERINSULINEMIA; CARDIOVASCULAR RISK;  
ELDERLY MEN)

95-3190 002 (INCREASED ABUNDANCE OF SPECIFIC SKELETAL-MUSCLE  
PROTEIN-TYROSINE PHOSPHATASES; ALPHA-B- **CRYSTALLIN** EXPRESSION)

95-0651 001 (CALCIUM CHANNELS; RAT CEREBELLAR GRANULE NEURONS; CA2+  
RELEASE)

95-3958 001 (RAT ISOLATED ANOCOCCYGEUS MUSCLE; SCORPION TOXINS;  
VOLTAGE-GATED ION CHANNELS; L- **ARGININE** -NITRIC OXIDE PATHWAY  
INVOLVEMENT; RELAXANT RESPONSES)

95-5343 001 (CAMP-DEPENDENT PROTEIN-KINASE; SIGNALING SPECIFICITY;  
PHOSPHATE PROTECTION)

Cited References:

ANDERSON EA, 1993, V21, P136, HYPERTENSION  
ANDERSON EA, 1991, V87, P2246, J CLIN INVEST  
BACKX PH, 1992, V257, P248, SCIENCE  
BERNE C, 1992, V35, P873, DIABETOLOGIA  
BRIGGS CA, 1981, V36, P1097, J NEUROCHEM  
CHANG SS, 1993, V20, P713, CLIN EXP PHARMACOL P  
CHENG JT, 1990, V190, P315, RES EXP MED  
DOOLEY DJ, 1987, V336, P467, N-S ARCH PHARMACOL  
DUNNETT CW, 1964, V20, P428, BIOMETRICS  
EDWARDS JG, 1989, V67, P2335, J APPL PHYSIOL  
HERTTING G, 1962, V18, P161, BRIT J PHARMACOL  
HOSKING DJ, 1978, V27, P1043, DIABETES  
HUANG WC, 1992, V218, P77, GENETIC HYPERTENSION  
HUNTER T, 1987, V50, P823, CELL  
JAIMETCHEVERRY G, 1980, V188, P513, BRAIN RES  
JONSSON G, 1981, V72, P173, EUR J PHARMACOL  
LAEMMLI UK, 1970, V227, P680, NATURE  
LAKOS Z, 1990, V1023, P41, BIOCHIM BIOPHYS ACTA  
LOWRY OH, 1951, V193, P265, J BIOL CHEM  
MERCER DW, 1978, V24, P480, CLIN CHEM  
MILLER RJ, 1987, V235, P46, SCIENCE  
MONDON CE, 1988, V37, P303, METABOLISM  
NACHSHEN DA, 1979, V16, P579, MOL PHARMACOL  
NARAHASHI T, 1974, V54, P813, PHYSIOL REV  
PATON WDM, 1968, V194, P13, J PHYSIOL  
REGAZZI R, 1990, V265, P15003, J BIOL CHEM  
RINK TJ, 1980, V595, P15, BIOCHIM BIOPHYS ACTA  
RITCHIE JM, 1977, V79, P1, REV PHYSIOL BIOCH PHAR  
RODRIGUEZPASCUAL F, 1995, V67, P149, NEUROSCIENCE  
ROGART RB, 1989, V86, P8170, P NATL ACAD SCI USA  
SAUTER A, 1983, V260, P330, BRAIN RES  
STARKE K, 1972, V195, P3309, ARCH INT PHARMACOD T  
SUZUKI K, 1976, V42, P399, J CLIN ENDOCR METAB  
SWISLOCKI A, 1990, V300, P104, AM J MED SCI  
TAGLIALATELA M, 1990, V1026, P126, BIOCHIM BIOPHYS ACTA  
TOWBIN H, 1979, V76, P4350, P NATL ACAD SCI USA  
WHITE MF, 1986, V17, P247, ENZYMES

...Abstract: a potential-sensitive dye.

chloride from the bathing medium. The induction of calcium ion influx by insulin into the synaptosomes...

...Research Fronts: MEN)

- 95-3190 002 (INCREASED ABUNDANCE OF SPECIFIC SKELETAL-MUSCLE PROTEIN-TYROSINE PHOSPHATASES; ALPHA-B- CRYSTALLIN EXPRESSION)
- 95-0651 001 (CALCIUM CHANNELS; RAT CEREBELLAR GRANULE NEURONS; CA2+ RELEASE)
- 95-3958 001 (RAT ISOLATED ANOCOCYGEUS MUSCLE; SCORPION TOXINS; VOLTAGE-GATED ION CHANNELS; L- ARGININE -NITRIC OXIDE PATHWAY INVOLVEMENT; RELAXANT RESPONSES)
- 95-5343 001 (CAMP-DEPENDENT PROTEIN-KINASE; SIGNALING SPECIFICITY...

5/9,K/4 (Item 4 from file: 34)

DIALOG(R) File 34:SciSearch(R) Cited Ref Sci

(c) 2007 The Thomson Corp. All rts. reserv.

05675361 Genuine Article#: WP266 Number of References: 62

**Title: The effect of hypotonicity, glutamine, and glycine on red cell preservation**

Author(s): Greenwalt TJ (REPRINT) ; Rugg N; Dumaswala UJ

Corporate Source: UNIV CINCINNATI, MED CTR, HOXWORTH BLOOD CTR, DEPT RES, 3131 HIGHLAND AVE, POB 670055/CINCINNATI//OH/45267 (REPRINT)

Journal: TRANSFUSION, 1997, V37, N3 (MAR), P269-276

ISSN: 0041-1132 Publication date: 19970300

Publisher: AMER ASSOC BLOOD BANKS, 8101 GLENBROOK RD, BETHESDA, MD 20814-2749

Language: English Document Type: ARTICLE

Geographic Location: USA

Subfile: CC LIFE--Current Contents, Life Sciences; CC CLIN--Current Contents, Clinical Medicine

Journal Subject Category: HEMATOLOGY

**Abstract:** BACKGROUND: Red cells (RBCs) stored in hypo-osmolar additive solutions with the same concentrations of adenine, dextrose, mannitol, and sodium chloride and varied amounts of ammonium, phosphate, glycerol, and glutamine were better preserved than RBCs in the standard additive solution (Adsol). Cell swelling occurred in all the experimental additives. This observation prompted the evaluation of glutamine and glycine alone, as well as a combination of glutamine and glycine, all of which have been described as producing swelling of rat liver cells.

**STUDY DESIGN AND METHODS:** Aliquots of RBCs were stored at 4 degrees C in Adsol or experimental additive solutions (EASs) all containing adenine, 2 mM; dextrose, 110 mM; mannitol, 55 mM; and sodium chloride, 50 mM. EAS 42 had, in addition, glutamine, 10 mM; glycine 5 mM; and phosphate, 20 mM. EAS 43 had glutamine, 10 mM; glycine, 10 mM; and phosphate 20 mM. EAS 44 had glutamine, 10 mM; EAS 45 had glutamine, 10 mM, and phosphate, 20 mM; and EAS 46 had only glycine, 10 mM. At intervals, measurements were made of mean corpuscular volume, mean corpuscular hemoglobin concentration, morphology, ATP, hemolysis, supernatant potassium, ammonia, pH, and microvesicles shed.

**RESULTS:** The initial mean corpuscular volumes were larger in all EASs than in Adsol, but the greatest difference was between EASs 44 and 46 (108 fL) and Adsol (86 fL) ( $p < 0.001$ ). The morphology scores were significantly better in all the EASs ( $p < 0.04$ ). The ATPs were

significantly greater in all the EASS ( $p < 0.001$ ), and highest in those with phosphate. Potassium leakage and hemolysis were less in the EASS ( $p < 0.001$ ). The ammonia levels were higher in all the EASS than in Adsol, with the exception of EAS 46. During storage, the extracorporeal and intracorporeal pH levels were essentially identical. The shedding of microvesicles was greatly reduced in all the EASS.

**CONCLUSION:** Cell swelling induced in RBCs after collection appears to improve preservation. Ammonia and phosphate enhance RBC ATP maintenance. Glycine decreases the formation of ammonia by RBCs stored in a hypotonic medium.

Identifiers--KeyWord Plus(R): AMINO-ACID-TRANSPORT; HUMAN-ERYTHROCYTES; KCL COTRANSPORT; MEMBRANE VESICULATION; STORED ERYTHROCYTES; ADDITIVE SOLUTION; CL COTRANSPORT; BLOOD-CELLS; RAT-LIVER; VOLUME

Research Fronts: 95-3190 001 (INCREASED ABUNDANCE OF SPECIFIC SKELETAL-MUSCLE PROTEIN-TYROSINE PHOSPHATASES; ALPHA-B- **CRYSTALLIN** EXPRESSION)

95-3483 001 (MULTIDRUG-RESISTANCE P-GLYCOPROTEIN; CELL VOLUME-ACTIVATED **CHLORIDE** CHANNELS; EXPRESSION PATTERN)

95-5062 001 (L- **ARGININE** TRANSPORT; NITRIC-OXIDE SYNTHASE ACTIVITY; SYSTEM Y(+))

#### Cited References:

- \*SAS STAT, 1988, SAS STAT V6 03
- ADAMS H, 1963, P539, METHOD ENZYMAT AN
- ANGELO S, 1994, V141, P183, J MEMBRANE BIOL
- BEUTLER E, 1987, V70, P1389, BLOOD
- BEUTLER E, 1990, P291, GLUTATHIONE METABOLI
- BEUTLER E, 1966, V6, P124, TRANSFUSION
- BEUTLER E, 1988, V28, P353, TRANSFUSION
- BRADFORD MM, 1976, V72, P248, ANAL BIOCHEM
- BRUGNARA C, 1988, V255, PC346, AM J PHYSIOL
- DANON D, 1964, V64, P668, J LAB CLIN MED
- DEBSKI B, 1981, V32, P73, ACTA PHYSIOL POL
- DEBSKI B, 1985, V36, P345, ACTA PHYSIOL POL
- DELOECKER WJ, 1975, V10, P157, CLIN ACTA
- DEUTICKE B, 1991, V1067, P111, BIOCHIM BIOPHYS ACTA
- DEVES R, 1992, V454, P491, J PHYSIOL-LONDON
- DUMASWALA UJ, 1994, V67, P139, VOX SANG
- DUMASWALA UJ, 1994, V67, P255, VOX SANG
- DUNHAM PB, 1990, P333, REGULATION POTASSIUM
- ELLORY JC, 1983, V42, P548, BIOMED BIOCHIM ACTA
- ELLORY JC, 1981, V320, P403, J PHYSIOL-LONDON
- ELLORY JC, 1983, V348, J PHYSIOL
- ENGLAND MD, 1986, V74, P134, CIRCULATION
- GERBER G, 1967, V18, P305, ACTA BIOL MED GER
- GOLDSTEIN S, 1984, V46, P725, INT J RADIAT BIOL
- GREENWALT TJ, 1984, V47, P261, VOX SANG
- GREENWALT TJ, 1993, V65, P87, VOX SANG
- GREENWALT TJ, 1996, V70, P6, VOX SANG
- HAUSSINGER D, 1991, V69, P1, BIOCH CELL BIOL
- HAUSSINGER D, 1990, V188, P689, EUR J BIOCHEM
- HENDERSON JF, 1979, V57, P873, CAN J BIOCHEM
- IMMERMAN KL, 1983, V20, P30, CRYOBIOLOGY
- JACOBASCH G, 1974, P55, CELLULAR MOL BIOL ER
- JENNINGS ML, 1990, V259, PC960, AM J PHYSIOL
- JOINER CH, 1993, V264, PC25, AM J PHYSIOL

KAJI DM, 1991, V260, PC176, AM J PHYSIOL  
 KLOPPICK E, 1967, V18, P37, J ACTA BIOL MED GER  
 LAUF PK, 1992, V263, PC917, AM J PHYSIOL  
 LAUF PK, 1985, V88, P1, J MEMBRANE BIOL  
 LUTZ HU, 1977, V73, P548, J CELL BIOL  
 MERYMAN HT, 1986, V26, P500, TRANSFUSION  
 MILLER GW, 1993, V53, P1211, LIFE SCI  
 MONDZAC A, 1965, V66, P526, J LAB CLIN MED  
 MOORE GL, 1981, V26, P167, BIOCHEM MED  
 OBERLEY LW, 1985, P417, HDB METHODS OXYGEN R  
 OLIVIERI O, 1993, V65, P95, VOX SANG  
 OSTLE B, 1975, P339, STAT RES BASCI CONCE  
 PARKER JC, 1989, P507, RED BLOOD CELL MEMBR  
 RAPOPORT SM, 1974, P93, CELLULAR MOL BIOL ER  
 RORTH M, 1972, V5, P1, SER HAEMATOL  
 ROSE ZB, 1970, V35, P177, ANAL BIOCHEM  
 SACHS JR, 1988, V92, P685, J GENERAL PHYSL  
 SARKADI B, 1991, V1071, P407, BIOCHIM BIOPHYS ACTA  
 TURNER S, 1987, V52, P177, VOX SANG  
 TURNER S, 1987, V52, P182, VOX SANG  
 TURNER S, 1987, V52, P186, VOX SANG  
 VINCENT N, 1992, V1175, P13, BIOCHIM BIOPHYS ACTA  
 WAGNER GM, 1987, V69, P1777, BLOOD  
 WAGNER GM, 1986, V108, P315, J LAB CLIN MED  
 WIDMANN FK, 1985, P64, TECHNICAL MANUAL  
 YOUNG JD, 1983, V216, P349, BIOCHEM J  
 YOUNG JD, 1982, P119, RED CELL MEMBRANES M  
 ZAR JH, 1974, P157, BIOSTAT ANAL

...Abstract: in hypo-osmolar additive solutions with the same concentrations of adenine, dextrose, mannitol, and sodium **chloride** and varied amounts of ammonium, phosphate, glycerol, and glutamine were better preserved than RBCs in...

...solutions (EASs) all containing adenine, 2 mM; dextrose, 110 mM; mannitol, 55 mM; and sodium **chloride**, 50 mM. EAS 42 had, in addition, glutamine, 10 mM; glycine 5 mM; and phosphate...

Research Fronts: 95-3190 001 (INCREASED ABUNDANCE OF SPECIFIC SKELETAL-MUSCLE PROTEIN-TYROSINE PHOSPHATASES; ALPHA-B- **CRYSTALLIN** EXPRESSION)

95-3483 001 (MULTIDRUG-RESISTANCE P-GLYCOPROTEIN; CELL VOLUME-ACTIVATED **CHLORIDE** CHANNELS; EXPRESSION PATTERN)

95-5062 001 (L- **ARGININE** TRANSPORT; NITRIC-OXIDE SYNTHASE ACTIVITY; SYSTEM Y(+))

5/9,K/5 (Item 5 from file: 34)

DIALOG(R)File 34:SciSearch(R) Cited Ref Sci

(c) 2007 The Thomson Corp. All rts. reserv.

05533867 Genuine Article#: WE933 Number of References: 41

Title: Inactivation and recovery of nitric oxide synthetic capability in cytokine-induced RAW 264.7 cells treated with 'irreversible' NO synthase inhibitors

Author(s): Wolff DJ (REPRINT) ; Lubeskie A; Li C

Corporate Source: UNIV MED & DENT NEW JERSEY, ROBERT WOOD JOHNSON MED SCH, DEPT PHARMACOL/PISCATAWAY//NJ/08854 (REPRINT)

Journal: ARCHIVES OF BIOCHEMISTRY AND BIOPHYSICS, 1997, V338, N1 (FEB 1), P

73-82

ISSN: 0003-9861 Publication date: 19970201

Publisher: ACADEMIC PRESS INC JNL-COMP SUBSCRIPTIONS, 525 B ST, STE 1900,  
SAN DIEGO, CA 92101-4495

Language: English Document Type: ARTICLE

Geographic Location: USA

Subfile: CC LIFE--Current Contents, Life Sciences

Journal Subject Category: BIOCHEMISTRY & MOLECULAR BIOLOGY; BIOPHYSICS

Abstract: As measured at 100  $\mu$ M extracellular **arginine**, aminoguanidine produced a time- and concentration-dependent inactivation of nitric oxide (NO) synthesis by cytokine-induced RAW cells. Inactivation obeyed first-order kinetics and occurred at a maximal rate of 0.22 min<sup>-1</sup> with a half-maximal inactivation rate observed at a concentration of 670  $\mu$ M aminoguanidine (K-I value). Inactivation of NO synthetic activity in the presence of N-G-methyl-L- **arginine** similarly followed first-order kinetics with a maximal inactivation rate of 0.07 min<sup>-1</sup> and a K-I value of 170  $\mu$ M. Inactivation of NO synthetic activity in the presence of diphenyliodonium **chloride** occurred with a maximal inactivation rate of 0.24 min<sup>-1</sup> with a K-I value of 14  $\mu$ M. Diphenyliodonium **chloride** also produced a first-order rate of inactivation of cytokine-inducible nitric oxide synthase (iNOS) activity affinity purified from cytokine-induced RAW cells with a maximal inactivation rate of its cytochrome c reductase activity of 0.24 min<sup>-1</sup> with a K-I value of 18  $\mu$ M. Cytokine-induced RAW cells were treated with aminoguanidine, N-G-methyl-L- **arginine**, and diphenyliodonium **chloride** at concentrations and for a time sufficient to completely inactivate NO synthesis by the cells and were allowed to recover in drug-free medium. Despite the presence of cycloheximide, NO synthetic rate recovered from 70 to 90% of its pretreatment activity over 4 h in cells exposed to either aminoguanidine or N-G-methyl-L-arginine but did not recover from exposure to diphenyliodonium **chloride**. Analysis by sucrose density gradient centrifugation of the cytochrome c reductase and citrulline-forming activities in extracts of cells recovered from aminoguanidine treatment revealed that recovery was accompanied by a diminished population of iNOS monomers with an increased population of iNOS dimers. This observation is consistent with the hypothesis that for the mechanism-based inactivator aminoguanidine, functional dimers can be assembled from "drug-undamaged" monomers during the recovery period.  
(C) 1997 Academic Press.

Descriptors--Author Keywords: aminoguanidine ; N-G-methyl-L- **arginine** ; diphenyliodonium **chloride** ; nitric oxide synthesis ; mechanism-based inactivation ; recovery ; intact cells

Identifiers--KeyWord Plus(R): METHYL-L- **ARGININE** ; IN-VIVO; AMINO GUANIDINE; MACROPHAGE; MECHANISM; DIPHENYLENEIODONIUM; IMIDAZOLE; REDUCTASE; ENZYME; POLYPEPTIDE

Research Fronts: 95-0388 003 (NITRIC-OXIDE SYNTHASE; ALDEHYDE FIXATION DIFFERENTIALLY AFFECTS DISTRIBUTION OF DIAPHORASE ACTIVITY; LIGHT-INDUCED FOS EXPRESSION)

95-1748 002 (INDUCIBLE NITRIC-OXIDE SYNTHASE; IN-VITRO ENDOTOXIN EXPOSURE INDUCES CONTRACTILE DYSFUNCTION IN ADULT-RAT CARDIAC MYOCYTES)

95-2212 001 (PEROXYNITRITE IN-VITRO; NITRIC-OXIDE SYNTHASE; HYDROXYL RADICAL; FORMATION OF 8-NITROGUANINE; PC12 CELLS)

95-3190 001 (INCREASED ABUNDANCE OF SPECIFIC SKELETAL-MUSCLE PROTEIN-TYROSINE PHOSPHATASES; ALPHA-B- **CRYSTALLIN** EXPRESSION)

Cited References:

ABUSOUD HM, 1995, V34, P11167, BIOCHEMISTRY-US

ALBAKRI QA, 1996, V271, P5414, J BIOL CHEM  
 BAEK KJ, 1993, V268, P21220, J BIOL CHEM  
 BRADFORD MM, 1976, V72, P248, ANAL BIOCHEM  
 BREDT DS, 1994, V63, P175, ANNU REV BIOCHEM  
 BREDT DS, 1994, V351, P714, NATURE  
 BREDT DS, 1990, V87, P682, P NATL ACAD SCI USA  
 CENDAN JC, 1996, V60, P284, J SURG RES  
 CROSS AH, 1994, V93, P2684, J CLIN INVEST  
 CROSS AR, 1986, V237, P111, BIOCHEM J  
 DEBELDER AJ, 1993, V341, P84, LANCET  
 FURFINE ES, 1994, V269, P26677, J BIOL CHEM  
 GRIFFITH OW, 1995, V57, P707, ANN REV PHYSL  
 HUANG Z, 1994, V265, P1883, SCIENCE  
 KITZ R, 1962, V237, P3245, J BIOL CHEM  
 KLATT P, 1992, V267, P11374, J BIOL CHEM  
 LAMAS S, 1992, V89, P6348, P NATL ACAD SCI USA  
 MARLETTA MA, 1994, V78, P927, CELL  
 MCCARTNEYFRANCI.N, 1993, V178, P749, J EXP MED  
 MONCADA S, 1995, V9, P1319, FASEB J  
 NARAYANAN K, 1995, V270, P11103, J BIOL CHEM  
 NATHAN C, 1992, V6, P3051, FASEB J  
 NOACK E, 1992, V1, P133, NEUROPROTOCOLS  
 OLKEN NM, 1991, V177, P828, BIOCHEM BIOPH RES CO  
 OLKEN NM, 1993, V27, P9677, BIOCHEMISTRY-US  
 OLKEN NM, 1994, V33, P14784, BIOCHEMISTRY-US  
 RAGAN CI, 1977, V163, P605, BIOCHEM J  
 RANDO RR, 1994, V36, P111, PHARMACOL REV  
 SCHMIDT HHH, 1992, V41, P615, MOL PHARMACOL  
 SILVERMAN RB, 1988, V1, P1, MECHANISM BASED ENZY  
 SIMMONS WW, 1996, V271, P11694, J BIOL CHEM  
 STUEHR DJ, 1991, V5, P98, FASEB J  
 STUHL MILLER DFE, 1995, V65, P68, J NEUROCHEM  
 TAMURA H, 1989, V105, P299, J BIOCHEM-TOKYO  
 TEW DG, 1993, V32, P10209, BIOCHEMISTRY-US  
 WANG YX, 1993, V110, P1232, BRIT J PHARMACOL  
 WOLFF DJ, 1994, V311, P293, ARCH BIOCHEM BIOPHYS  
 WOLFF DJ, 1994, V314, P360, ARCH BIOCHEM BIOPHYS  
 WOLFF DJ, 1995, V316, P290, ARCH BIOCHEM BIOPHYS  
 WOLFF DJ, 1992, V285, P201, BIOCHEM J  
 WOLFF DJ, 1993, V268, P9425, J BIOL CHEM

Abstract: As measured at 100  $\mu$  M extracellular **arginine**, aminoguanidine produced a time- and concentration-dependent inactivation of nitric oxide (NO) synthesis by cytokine...

...I value). Inactivation of NO synthetic activity in the presence of N-G-methyl-L- **arginine** similarly followed first-order kinetics with a maximal inactivation rate of 0.07 min<sup>-1</sup>...

...value of 170  $\mu$  M. Inactivation of NO synthetic activity in the presence of diphenyliodonium **chloride** occurred with a maximal inactivation rate of 0.24 min<sup>-1</sup> with a K-I value of 14  $\mu$  M. Diphenyliodonium **chloride** also produced a first-order rate of inactivation of cytokine-inducible nitric oxide synthase (iNOS...

...18  $\mu$  M. Cytokine-induced RAW cells were treated with aminoguanidine, N-G-methyl-L- **arginine**, and diphenyliodonium **chloride** at concentrations and for a time sufficient to completely inactivate NO



synthesis by the cells...

...aminoguanidine or N-G-methyl-L-arsnine but did not recover from exposure to diphenyliodonium chloride . Analysis by sucrose density gradient centrifugation of the cytochrome c reductase and citrulline-forming activities...

...Identifiers--METHYL-L- ARGININE ; IN-VIVO; AMINOGUANIDINE; MACROPHAGE; MECHANISM; DIPHENYLENEIODONIUM; IMIDAZOLE; REDUCTASE; ENZYME; POLYPEPTIDE

...Research Fronts: CELLS)

95-3190 001 (INCREASED ABUNDANCE OF SPECIFIC SKELETAL-MUSCLE PROTEIN-TYROSINE PHOSPHATASES; ALPHA-B- CRYSTALLIN EXPRESSION)

5/9,K/6 (Item 6 from file: 34)

DIALOG(R) File 34:SciSearch(R) Cited Ref Sci

(c) 2007 The Thomson Corp. All rts. reserv.

05463541 Genuine Article#: WA564 Number of References: 51

Title: **EXPRESSION OF THE CYSTIC-FIBROSIS PHENOTYPE IN A RENAL AMPHIBIAN EPITHELIAL-CELL LINE**

Author(s): LING BN; ZUCKERMAN JB; LIN CM; HARTE BJ; MCNULTY KA; SMITH PR; GOMEZ LM; WORRELL RT; EATON DC; KLEYMAN TR

Corporate Source: DEPT VET AFFAIRS, MED CTR, UNIV & WOODLAND

AVE/PHILADELPHIA//PA/19104; DEPT VET AFFAIRS, MED

CTR/PHILADELPHIA//PA/19104; DEPT VET AFFAIRS MED CTR/ATLANTA//GA/30322;

EMORY UNIV, DIV RENAL/ATLANTA//GA/30322; EMORY UNIV, DEPT

MED/ATLANTA//GA/30322; EMORY UNIV, DEPT PHYSIOL/ATLANTA//GA/30322; EMORY

UNIV, CTR CELL & MOL SIGNALING/ATLANTA//GA/30322; UNIV PENN, DEPT

MED/PHILADELPHIA//PA/19104; UNIV PENN, DEPT

PHYSIOL/PHILADELPHIA//PA/19104; ALLEGHENY UNIV HLTH SCI, DEPT

PHYSIOL/PHILADELPHIA//PA/19129

Journal: JOURNAL OF BIOLOGICAL CHEMISTRY, 1997, V272, N1 (JAN 3), P594-600

ISSN: 0021-9258

Language: ENGLISH Document Type: ARTICLE

Geographic Location: USA

Subfile: Science Citation Index; SciSearch; CC LIFE--Current Contents, Life Sciences

Journal Subject Category: BIOCHEMISTRY & MOLECULAR BIOLOGY

Abstract: Mutations in a Cl<sup>-</sup> channel (cystic fibrosis transmembrane conductance regulator or CFTR) are responsible for the cystic fibrosis (CF) phenotype, Increased Na<sup>+</sup> transport rates are observed in CF airway epithelium, and recent studies suggest that this is due to an increase in Na<sup>+</sup> channel open probability (P<sub>o</sub>). The Xenopus renal epithelial cell line, A6, expresses both cAMP-activated 8-picosiemen (pS) Cl<sup>-</sup> channels and amiloride-sensitive 4-pS Na<sup>+</sup> channels, and provides a model system for examining the interactions of CFTR and epithelial Na<sup>+</sup> channels. A6 cells express CFTR mRNA, as demonstrated by reverse transcriptase-polymerase chain reaction and partial sequence analysis. A phosphorothioate antisense oligonucleotide, complementary to the 5' end of the open reading frame of Xenopus CFTR, was used to inhibit functional expression of CFTR in A6 cells. Parallel studies utilized the corresponding sense oligonucleotide as a control. CFTR protein expression was markedly reduced in cells incubated with the antisense oligonucleotide. Incubation of A6 cells with the antisense oligonucleotide led to inhibition of forskolin-activated amiloride-insensitive short circuit current (I<sub>sc</sub>). After a 30-min

exposure to 10  $\mu$  M forskolin, 8-pS Cl<sup>-</sup> channel activity was detected in only 1 of 31 (3%) cell-attached patches on cells treated with antisense oligonucleotide, compared to 5 of 19 (26%) patches from control cells. A shift in the single-channel current-voltage relationship derived from antisense-treated cells was also consistent with a reduction in Cl<sup>-</sup> reabsorption. Both amiloride-sensitive I<sub>sc</sub> and Na<sup>+</sup> channel P<sub>o</sub> were significantly increased in antisense-treated, forskolin-stimulated A6 cells, when compared with forskolin-stimulated controls. These data suggest that the regulation of Na<sup>+</sup> channels by CFTR is not limited to respiratory epithelia and to epithelial cells in culture overexpressing CFTR and epithelial Na<sup>+</sup> channels.

Identifiers--Keywords Plus: TRANSMEMBRANE CONDUCTANCE REGULATOR; PROTEIN-KINASE-C; NA<sup>+</sup> CHANNELS; SODIUM-CHANNELS; CHLORIDE CHANNELS; ANTISENSE OLIGODEOXYNUCLEOTIDE; ARGININE -VASOPRESSIN; AIRWAY EPITHELIA; CFTR; A6

Research Fronts: 95-0327 004 (CYSTIC-FIBROSIS TRANSMEMBRANE CONDUCTANCE REGULATOR GENE; DIETARY-CHANGES IMPROVE SURVIVAL OF CFTR S489X HOMOZYGOUS MUTANT MOUSE)

95-5061 002 (STRUCTURAL GENE; GLTC-DEPENDENT REGULATION OF BACILLUS-SUBTILIS GLUTAMATE SYNTHASE EXPRESSION; ARABIDOPSIS TYPE-1 PROTEIN PHOSPHATASE)

95-3190 001 (INCREASED ABUNDANCE OF SPECIFIC SKELETAL-MUSCLE PROTEIN-TYROSINE PHOSPHATASES; ALPHA-B- CRYSTALLIN EXPRESSION)

95-4481 001 (K<sup>+</sup> CHANNELS IN CULTURED RAT NEURONAL CELLS; DIFFERENT GATING KINETICS; SINGLE NMDA RECEPTOR CURRENTS; CARDIAC SARCOPLASMIC-RETICULUM)

#### Cited References:

- ANDERSON MP, 1991, V253, P202, SCIENCE  
BAO HF, 1996, V10, A137, FASEB J  
BENOS DJ, 1995, V143, P1, J MEMBRANE BIOL  
BERGER HA, 1991, V88, P1422, J CLIN INVEST  
BOUCHER RC, 1991, V290, P105, ADV EXP MED BIOL  
BOUCHER RC, 1986, V78, P1245, J CLIN INVEST  
BOUCHER RC, 1988, V405, P77, J PHYSIOL-LONDON  
BRADBURY NA, 1992, V256, P530, SCIENCE  
CHALFANT ML, 1993, V264, P1480, AM J PHYSIOL  
CHINET TC, 1994, V266, P1061, AM J PHYSIOL  
CHOMCZYNSKI P, 1987, V162, P156, ANAL BIOCHEM  
COLQUHOUN D, 1983, P191, SINGLE CHANNEL RECOR  
COUPAYEGGERARD B, 1996, V271, P1639, AM J PHYSIOL  
DENNING GM, 1992, V358, P761, NATURE  
DISSER J, 1991, V290, P133, ADV EXP MED BIOL  
DUSZYK M, 1989, V67, P1362, CAN J PHYSIOL PHARM  
EATON DC, 1995, V48, P941, KIDNEY INT  
EGAN M, 1992, V358, P581, NATURE  
GALLEGO MS, 1996, V271, F451, AM J PHYSIOL  
GREGORY RJ, 1990, V347, P382, NATURE  
GRUBB BR, 1994, V266, P1478, AM J PHYSIOL  
JOHNSON LG, 1995, V95, P1377, J CLIN INVEST  
JOVOV B, 1995, V270, P1521, J BIOL CHEM  
JOVOV B, 1995, V270, P9194, J BIOL CHEM  
KEMENDY AE, 1992, V263, C825, AM J PHYSIOL  
KLEYMAN TR, 1994, V266, F506, AM J PHYSIOL  
KOKKO KE, 1994, V267, P1414, AM J PHYSIOL  
KOPELMAN H, 1993, V91, P1253, J CLIN INVEST  
KRAUSS RD, 1992, V263, P1147, AM J PHYSIOL  
LING BN, 1989, V256, P1094, AM J PHYSIOL

LING BN, 1992, V90, P1328, J CLIN INVEST  
 LING BN, 1994, V93, P829, J CLIN INVEST  
 MA H, 1996, V270, F798, AM J PHYSIOL  
 MA H, 1996, V9, P1284, J AM SOC NEPHROL  
 MARUNAKA Y, 1990, V258, C352, AM J PHYSIOL  
 MARUNAKA Y, 1991, V258, C352, AM J PHYSIOL  
 OH YS, 1993, V265, C85, AM J PHYSIOL  
 PUOTI A, 1995, V269, C188, AM J PHYSIOL  
 RIORDAN JR, 1993, V55, P609, ANN REV PHYSIOL  
 RIORDAN JR, 1989, V245, P1066, SCIENCE  
 SAMBROOK J, 1989, MOL CLONING LAB MANU  
 SANGER F, 1977, V74, P5463, P NATL ACAD SCI USA  
 SCHLANGER LE, 1994, V45, P1070, KIDNEY INT  
 SCHREIBER M, 1996, V49, P82, KIDNEY INT  
 SCHWIEBERT EM, 1995, V81, P1063, CELL  
 SMITH JJ, 1994, V93, P1307, J CLIN INVEST  
 SORSCHER EJ, 1991, V88, P7759, P NATL ACAD SCI USA  
 STUTTS MJ, 1995, V269, P847, SCIENCE  
 TOWBIN H, 1979, V76, P4350, P NATL ACAD SCI USA  
 TUCKER SJ, 1992, V1, P77, HUM MOL GENET  
 WAGNER JA, 1992, V89, P6785, P NATL ACAD SCI USA

...Identifiers--TRANSMEMBRANE CONDUCTANCE REGULATOR; PROTEIN-KINASE-C; NA+ CHANNELS; SODIUM-CHANNELS; **CHLORIDE** CHANNELS; ANTISENSE OLIGODEOXYNUCLEOTIDE; **ARGININE** -VASOPRESSIN; AIRWAY EPITHELIA; CFTR; A6

...Research Fronts: PHOSPHATASE)

95-3190 001 (INCREASED ABUNDANCE OF SPECIFIC SKELETAL-MUSCLE

PROTEIN-TYROSINE PHOSPHATASES; ALPHA-B- **CRYSTALLIN** EXPRESSION)

95-4481 001 (K+ CHANNELS IN CULTURED RAT NEURONAL CELLS; DIFFERENT GATING KINETICS; SINGLE...

5/9,K/7 (Item 7 from file: 34)

DIALOG(R)File 34:SciSearch(R) Cited Ref Sci

(c) 2007 The Thomson Corp. All rts. reserv.

05457800 Genuine Article#: WA251 Number of References: 31

Title: **AMYLIN AND FOOD-INTAKE IN MICE - EFFECTS ON MOTIVATION TO EAT AND MECHANISM OF ACTION**

Author(s): MORLEY JE; SUAREZ MD; MATTAMAL M; FLOOD JF

Corporate Source: ST LOUIS UNIV,SCH MED M239,DIV GERIATR MED,1402S GRAND

BLVD/ST LOUIS//MO/63104; VET ADM MED CTR,CTR GERIATR RES EDUC & CLIN/ST LOUIS//MO/63106

Journal: PHARMACOLOGY BIOCHEMISTRY AND BEHAVIOR, 1997, V56, N1 (JAN), P 123-129

ISSN: 0091-3057

Language: ENGLISH Document Type: ARTICLE

Geographic Location: USA

Subfile: Science Citation Index; SciSearch; CC LIFE--Current Contents, Life Sciences

Journal Subject Category: PHARMACOLOGY & PHARMACY

Abstract: Amylin is a hormone produced by the pancreatic islets of Langerhans. Amylin decreased food pellet consumption. Amylin also decreased lever pressing for milk solution whether or not the mice were prefed. Amylin did not produce a conditioned taste aversion in a two bottle test, whereas lithium chloride did. In addition, L- arginine ,

a precursor for nitric oxide synthesis, was demonstrated to inhibit the ability of amylin to decrease food intake. Amylin did not alter nitric oxide synthase activity in the fundus of the stomach. These studies demonstrated that amylin inhibits food intake at a higher range of doses than is typical of anorectic agents such as cholecystokinin. Amylin does not appear to decrease food intake by reducing the release of nitric oxide but may affect appetite by modulating serum glucose levels when co-released with insulin. Copyright (C) 1997 Elsevier

Science Inc.

Descriptors--Author Keywords: AMYLIN ; APPETITE ; FOOD INTAKE ; NITRIC OXIDE ; L- ARGININE ; NITRIC OXIDE SYNTHASE ; LEVER PRESS ; ANOREXIA ; MOTIVATION

Identifiers--KeyWords Plus: GUT PEPTIDES; RATS; SATIETY; MODULATION; INJECTION; APPETITE; WEIGHT

Research Fronts: 95-3190 001 (INCREASED ABUNDANCE OF SPECIFIC SKELETAL-MUSCLE PROTEIN-TYROSINE PHOSPHATASES; ALPHA-B- CRYSTALLIN EXPRESSION)

95-4861 001 (LICKING BEHAVIOR IN RATS; GASTRIN-RELEASING PEPTIDE; INGESTIVE TASTE REACTIVITY)

95-8036 001 (OBESE ZUCKER RATS; CENTRAL INSULIN; HYPOTHALAMIC PARAVENTRICULAR NUCLEUS; INVOLVEMENT OF NEUROPEPTIDE-Y; OPIOID ANTAGONIST NALOXONE)

#### Cited References:

BALASUBRAMANIAM A, 1991, V12, P919, PEPTIDES  
BILLINGTON CJ, 1983, V245, R920, AM J PHYSIOL  
BRADFORD MM, 1976, V72, P248, ANAL BIOCHEM  
CALIGNANO A, 1993, V231, P415, EUR J PHARMACOL  
CAREY RJ, 1973, V1, P227, PHARM BIO B  
CHANCE WT, 1991, V539, P352, BRAIN RES  
CHANCE WT, 1993, V607, P185, BRAIN RES  
CHANCE WT, 1992, V13, P961, PEPTIDES  
DAVIS JL, 1977, V19, P55, BEHAV NEURAL BIOL  
DESAI KM, 1991, V88, P1490, P NATL ACAD SCI USA  
DEUTSCH JA, 1977, V266, P196, NATURE  
EDWARDS BJA, 1992, V51, P1899, LIFE SCI  
FLOOD JF, 1987, V421, P280, BRAIN RES  
FLOOD JF, 1990, V11, P265, PEPTIDES  
GIBBS J, 1979, V282, P208, NATURE  
KEPPEL G, 1973, DESIGN ANAL RESEARCH  
KRAHN DD, 1984, V5, P861, PEPTIDES  
LESAUTER J, 1987, V253, R719, AM J PHYSIOL  
MONCADA S, 1992, V145, P201, ACTA PHYSIOL SCAND  
MORLEY JE, 1994, V267, R178, AM J PHYSIOL  
MORLEY JE, 1990, V10, P383, ANNU REV NUTR  
MORLEY JE, 1987, V8, P256, ENDOCR REV  
MORLEY JE, 1991, V49, P707, LIFE SCI  
MORLEY JE, 1992, V51, P1285, LIFE SCI  
MORLEY JE, 1991, V12, P865, PEPTIDES  
MORLEY JE, 1983, V19, P577, PHARMACOL BIOCHEM BE  
MORLEY JE, 1993, V44, P577, PHARMACOL BIOCHEM BE  
NACHMAN M, 1973, V10, P73, PHYSIOL BEHAV  
SMITH GP, 1984, V43, P2889, FED PROC  
SQUADRITO F, 1994, V33, P83, NEUROPHARMACOLOGY  
WINER BJ, 1971, STATISTICAL PRINCIPAL

...Abstract: Amylin did not produce a conditioned taste aversion in a two

bottle test, whereas lithium **chloride** did. In addition, L- **arginine** , a precursor for nitric oxide synthesis, was demonstrated to inhibit the ability of amylin to...

Research Fronts: 95-3190 001 (INCREASED ABUNDANCE OF SPECIFIC SKELETAL-MUSCLE PROTEIN-TYROSINE PHOSPHATASES; ALPHA-B- **CRYSTALLIN** EXPRESSION)

95-4861 001 (LICKING BEHAVIOR IN RATS; GASTRIN-RELEASING PEPTIDE; INGESTIVE TASTE REACTIVITY)

95...

5/9,K/8 (Item 8 from file: 34)

DIALOG(R)File 34:SciSearch(R) Cited Ref Sci

(c) 2007 The Thomson Corp. All rts. reserv.

00857477 Genuine Article#: FC075 Number of References: 22

Title: INTERACTIONS BETWEEN ION-EXCHANGE AND METABOLISM IN ERYTHROCYTES OF THE RAINBOW-TROUT ONCORHYNCHUS-MYKISS

Author(s): TUFTS BL; BOUTILIER RG

Corporate Source: DALHOUSIE UNIV,DEPT BIOL/HALIFAX B3H 4J1/NS/CANADA/

Journal: JOURNAL OF EXPERIMENTAL BIOLOGY, 1991, V156, MAR, P139-151

Language: ENGLISH Document Type: ARTICLE

Geographic Location: CANADA

Subfile: SciSearch; CC LIFE--Current Contents, Life Sciences; CC AGRI--

Current Contents, Agriculture, Biology & Environmental Sciences

Journal Subject Category: BIOLOGY

Abstract: Experiments were carried out to investigate the relationship between ion exchange and energy metabolism in rainbow trout erythrocytes in vitro. Under resting conditions, the sodium/potassium pump accounts for 20% of the cellular energy budget. In the presence of the beta-adrenergic agonist isoproterenol, however, this increases to 43%. Inhibition of the sodium/potassium pump with ouabain results in greater increases in erythrocyte water content and sodium and **chloride** concentrations and a greater decrease in erythrocyte potassium concentration following stimulation by isoproterenol. Moreover, the decrease in erythrocyte NTP levels observed following adrenergic stimulation does not occur when the sodium/potassium pump is inhibited with ouabain. Inhibition of the sodium/potassium pump also abolishes the increase in oxygen consumption by the cells which normally takes place following adrenergic stimulation. Finally, depletion of erythrocyte NTP levels by the sodium ionophore monensin or by previous incubation with nitrogen does not result in a significant increase in oxygen consumption. Thus, catecholamines appear to be crucial for the metabolic-membrane coupling that occurs following adrenergic stimulation in rainbow trout erythrocytes.

Descriptors--Author Keywords: ERYTHROCYTES; TROUT; ION EXCHANGE; METABOLISM; ONCORHYNCHUS-MYKISS

Identifiers--KeyWords Plus: RED-CELLS; PROTEIN-PHOSPHORYLATION; ADRENERGIC-STIMULATION; FISH ERYTHROCYTES; HORMONAL-CONTROL; PH REGULATION; MEMBRANE; TRANSPORT; VOLUME; INVIVO

Research Fronts: 89-1358 001 (RAINBOW-TROUT (SALMO-GAIRDNERI); ACID-BASE REGULATION FOLLOWING EXHAUSTIVE EXERCISE; MARINE FISH; RESPIRATORY ADAPTATIONS; ENZYMES OF **ARGININE** METABOLISM)

89-4150 001 (PHOSPHORYLATION OF PROTEINS; CATALYTIC SUBUNIT; GLYCOGEN-SYNTHASE ACTIVITY; CASEIN KINASE-2; LENS ALPHA- **CRYSTALLIN** A-CHAIN; PHOSPHODIESTER LINKAGE)

Cited References:

ALBERS C, 1985, V61, P209, RESP PHYSIOL  
BARNES DM, 1986, V234, P286, SCIENCE  
BORGESSE F, 1986, V87, P551, J GEN PHYSIOL  
BORGESSE F, 1987, V382, P145, J PHYSIOL-LONDON  
BOURNE PK, 1982, V101, P93, J EXP BIOL  
BOUTILIER RG, 1989, V67, P2986, CAN J ZOOL  
COHEN P, 1985, V151, P439, EUR J BIOCHEM  
COHEN P, 1982, V296, P613, NATURE  
COSSINS A, 1989, V340, P20, NATURE  
DEVRIES AL, 1982, V324, P51, J PHYSIOL-LONDON  
FERGUSON RA, 1989, V143, P149, J EXP BIOL  
FERGUSON RA, 1989, V143, P133, J EXP BIOL  
FERGUSON RA, 1988, V74, P65, RESPIR PHYSIOL  
HOFFMANN EK, 1989, V69, P315, PHYSIOL REV  
KALMAN M, 1984, V56, P539, EXP BRAIN RES  
MAHE Y, 1985, V116, P199, EUR J PHARMACOL  
MAIRBAURL H, 1981, P311, RED CELL  
NIKINMAA M, 1990, V67, P3039, CAN J ZOOL  
NIKINMAA M, 1983, V152, P67, J COMP PHYSIOL  
RAPOPORT SM, 1985, P333, CIRCULATION RESPIRAT  
SMITH LS, 1964, V21, P711, J FISH RES BOARD CAN  
TUCKER VA, 1967, V23, P410, J APPL PHYSIOL

...Abstract: potassium pump with ouabain results in greater increases in erythrocyte water content and sodium and chloride concentrations and a greater decrease in erythrocyte potassium concentration following stimulation by isoproterenol. Moreover, the...

...Research Fronts: TROUT (SALMO-GAIRDNERI); ACID-BASE REGULATION FOLLOWING EXHAUSTIVE EXERCISE; MARINE FISH; RESPIRATORY ADAPTATIONS; ENZYMES OF ARGININE METABOLISM)

89-4150 001 (PHOSPHORYLATION OF PROTEINS; CATALYTIC SUBUNIT;  
GLYCOGEN-SYNTHASE ACTIVITY; CASEIN KINASE-2; LENS ALPHA- CRYSTALLIN  
A-CHAIN; PHOSPHODIESTER LINKAGE)

5/9,K/9 (Item 1 from file: 73)

DIALOG(R)File 73:EMBASE

(c) 2007 Elsevier B.V. All rts. reserv.

13167229 EMBASE No: 2005234088

**Alexander-disease mutation of GFAP causes filament disorganization and decreased solubility of GFAP**

Hsiao V.C.; Tian R.; Long H.; Perng M.D.; Brenner M.; Quinlan R.A.;  
Goldman J.E.

J.E. Goldman, Department of Pathology, Center for Neurobiology and  
Behavior, Columbia University, New York, NY 10032 United States

AUTHOR EMAIL: jeg5@columbia.edu

Journal of Cell Science ( J. CELL SCI. ) (United Kingdom) 01 MAY 2005,  
118/9 (2057-2065)

CODEN: JNCSA ISSN: 0021-9533

DOCUMENT TYPE: Journal ; Article

LANGUAGE: ENGLISH SUMMARY LANGUAGE: ENGLISH

NUMBER OF REFERENCES: 66

Alexander disease is a fatal neurological illness characterized by white-matter degeneration and the formation of astrocytic cytoplasmic inclusions called Rosenthal fibers, which contain the intermediate filament

glial fibrillary acidic protein (GFAP), the small heat-shock proteins HSP27 and alphaB- **crystallin** , and ubiquitin. Many Alexander-disease patients are heterozygous for one of a set of point mutations in the GFAP gene, all of which result in amino acid substitutions. The biological effects of the most common alteration, R239C, were tested by expressing the mutated protein in cultured cells by transient transfection. In primary rat astrocytes and Cos-7 cells, the mutant GFAP was incorporated into filament networks along with the endogenous GFAP and vimentin, respectively. In SW13VimSUP- cells, which have no endogenous cytoplasmic intermediate filaments, wild-type human GFAP frequently formed filamentous bundles, whereas the R239C GFAP formed 'diffuse' and irregular patterns. Filamentous bundles of R239C GFAP were sometimes formed in SW13VimSUP- cells when wild-type GFAP was co-transfected. Although the presence of a suitable coassembly partner (vimentin or GFAP) reduced the potential negative effects of the R239C mutation on GFAP network formation, the mutation affected the stability of GFAP in cells in a dominant fashion. Extraction of transfected SW13VimSUP- cells with Triton-X-100-containing buffers showed that the mutant GFAP was more resistant to solubilization at elevated KCl concentrations. Both wild-type and R239C GFAP assembled into 10 nm filaments with similar morphology in vitro. Thus, although the R239C mutation does not appear to affect filament formation per se, the mutation alters the normal solubility and organization of GFAP networks.

#### DRUG DESCRIPTORS:

\*glial fibrillary acidic protein--endogenous compound--ec  
heat shock protein 27--endogenous compound--ec; alpha **crystallin**  
--endogenous compound--ec; beta **crystallin** --endogenous compound--ec;  
ubiquitin--endogenous compound--ec; amino acid--endogenous compound--ec;  
mutant protein--endogenous compound--ec; vimentin--endogenous compound--ec;  
triton x 100; buffer; potassium **chloride** ; **arginine** --endogenous compound  
--ec; cysteine--endogenous compound--ec

#### MEDICAL DESCRIPTORS:

\*Alexander disease--diagnosis--di; \*Alexander disease--etiology--et  
gene mutation; neurologic disease--etiology--et; clinical feature; white  
matter; astrocyte; cell inclusion; heterozygosity; point mutation; amino  
acid substitution; protein expression; cell culture; genetic transfection;  
cytoplasm; wild type; protein assembly; intermediate filament; protein  
stability; solubilization; concentration (parameters); cell structure; in  
vitro study; protein structure; human; nonhuman; rat; controlled study;  
human cell; animal cell; article; priority journal

CAS REGISTRY NO.: 60267-61-0 (ubiquitin); 65072-01-7 (amino acid);  
7447-40-7 (potassium **chloride** ); 1119-34-2, 15595-35-4, 7004-12-8,  
74-79-3 ( **arginine** ); 4371-52-2, 52-89-1, 52-90-4 (cysteine)

#### SECTION HEADINGS:

- 005 General Pathology and Pathological Anatomy
- 008 Neurology and Neurosurgery
- 022 Human Genetics
- 029 Clinical and Experimental Biochemistry

...intermediate filament glial fibrillary acidic protein (GFAP), the  
small heat-shock proteins HSP27 and alphaB- **crystallin** , and ubiquitin.  
Many Alexander-disease patients are heterozygous for one of a set of point  
...

#### DRUG DESCRIPTORS:

heat shock protein 27--endogenous compound--ec; alpha **crystallin**  
--endogenous compound--ec; beta **crystallin** --endogenous compound--ec;  
ubiquitin--endogenous compound--ec; amino acid--endogenous compound--ec;

mutant protein--endogenous compound--ec; vimentin--endogenous compound--ec;  
triton x 100; buffer; potassium **chloride** ; **arginine** --endogenous compound  
--ec; cysteine--endogenous compound--ec  
CAS REGISTRY NO.: 60267-61-0 (ubiquitin); 65072-01-7 (amino acid);  
7447-40-7 (potassium **chloride** ); 1119-34-2...

...74-79-3 ( **arginine** ); 4371-52-2...

5/9,K/10 (Item 2 from file: 73)  
DIALOG(R) File 73:EMBASE  
(c) 2007 Elsevier B.V. All rts. reserv.

07461271 EMBASE No: 1998370560

**Human procarboxypeptidase U, or thrombin-activable fibrinolysis inhibitor, is a substrate for transglutaminases: Evidence for transglutaminase-catalyzed cross-linking to fibrin**

Valnickova Z.; Enghild J.J.

J.J. Enghild, Box 3712, Duke University Medical Center, Durham, NC 27710  
United States

AUTHOR EMAIL: enghi001@mc.duke.edu

Journal of Biological Chemistry ( J. BIOL. CHEM. ) (United States) 16

OCT 1998, 273/42 (27220-27224)

CODEN: JBCHA ISSN: 0021-9258

DOCUMENT TYPE: Journal; Article

LANGUAGE: ENGLISH SUMMARY LANGUAGE: ENGLISH

NUMBER OF REFERENCES: 45

Procarboxypeptidase U (EC 3.4.17.20) (pro-CpU), also known as plasma procarboxypeptidase B and thrombin-activable fibrinolysis inhibitor, is a human plasma protein that has been implicated in the regulation of fibrinolysis. In this study, we show that pro-CpU serves as a substrate for transglutaminases. Both factor XIIIa and tissue transglutaminase catalyzed the polymerization of pro-CpU and the cross-linking to fibrin as well as the incorporation of 5-dimethylaminonaphthalene-1-sulfonyl cadaverine (dansylcadaverine), [sup 1sup 4C]putrescine, and dansyl-PGGQQIV. These findings show that pro-CpU contains both amine acceptor (Gln) and amine donor (Lys) residues. The amine acceptor residues were identified as Glnsup 2, Glnsup 5, and Glnsup 2sup 9sup 2, suggesting that both the activation peptide and the mature enzyme participate in the cross-linking reaction. These observations imply that transglutaminases may mediate covalent binding of pro-CpU to other proteins and cell surfaces in vivo. In particular, factor XIIIa may cross-link pro-CpU to fibrin during the latter part of the coagulation cascade, thereby helping protect the newly formed fibrin clot from premature plasmin degradation. Moreover, the cross-linking may facilitate the activation of pro-CpU, stabilize the enzymatic activity, and protect the active enzyme from further degradation.

#### DRUG DESCRIPTORS:

\*protein glutamine gamma glutamyltransferase--endogenous compound--ec; \*  
antifibrinolytic agent--endogenous compound--ec  
dansylcadaverine--endogenous compound--ec; putrescine--endogenous compound  
--ec; dansyl **chloride** --endogenous compound--ec; glutamine--endogenous  
compound--ec; lysine--endogenous compound--ec; blood clotting factor 13  
--endogenous compound--ec; fibrin--endogenous compound--ec; plasmin  
--endogenous compound--ec; **arginine** --endogenous compound--ec; alpha 2  
antiplasmin--endogenous compound--ec; amine--endogenous compound--ec;



plasminogen activator inhibitor 1--endogenous compound--ec; beta  
**crystallin** --endogenous compound--ec; fibrinogen--endogenous compound--ec;  
vitronectin--endogenous compound--ec; unclassified drug

MEDICAL DESCRIPTORS:

\*fibrinolysis; \*enzyme substrate

fibrin polymerization; covalent bond; fibrin clot; enzyme activity; enzyme  
stability; enzyme degradation; liver; sequence homology; carboxy terminal  
sequence; enzyme active site; protein cross linking; enzyme specificity;  
human; nonhuman; human tissue; animal tissue; article; priority journal

DRUG TERMS (UNCONTROLLED): procarboxypapetidase u--endogenous compound--ec;  
thrombin activable fibrinolysis inhibitor

CAS REGISTRY NO.: 80146-85-6 (protein glutamine gamma glutamyltransferase);  
10121-91-2 (dansylcadaverine); 110-60-1, 333-93-7 (putrescine);  
605-65-2 (dansyl **chloride** ); 56-85-9, 6899-04-3 (glutamine); 56-87-1,  
6899-06-5, 70-54-2 (lysine); 9013-56-3 (blood clotting factor 13);  
9001-31-4 (fibrin); 9001-90-5, 9004-09-5 (plasmin); 1119-34-2,  
15595-35-4, 7004-12-8, 74-79-3 ( **arginine** ); 140208-23-7 (plasminogen  
activator inhibitor 1); 9001-32-5 (fibrinogen)

SECTION HEADINGS:

025 Hematology

029 Clinical and Experimental Biochemistry

DRUG DESCRIPTORS:

dansylcadaverine--endogenous compound--ec; putrescine--endogenous compound  
--ec; dansyl **chloride** --endogenous compound--ec; glutamine--endogenous  
compound--ec; lysine--endogenous compound--ec; blood clotting factor 13  
--endogenous compound--ec; fibrin--endogenous compound--ec; plasmin  
--endogenous compound--ec; **arginine** --endogenous compound--ec; alpha 2  
antiplasmin--endogenous compound--ec; amine--endogenous compound--ec;  
plasminogen activator inhibitor 1--endogenous compound--ec; beta  
**crystallin** --endogenous compound--ec; fibrinogen--endogenous compound--ec;  
vitronectin--endogenous compound--ec; unclassified drug

...CAS REGISTRY NO.: 333-93-7 (putrescine); 605-65-2 (dansyl **chloride** );  
56-85-9...

...74-79-3 ( **arginine** ); 140208-23-7 (plasminogen activator inhibitor 1);  
9001-32-5 (fibrinogen)

? s arginine(n)hydrochloride

416125 ARGININE

382785 HYDROCHLORIDE

S6 942 ARGININE(N)HYDROCHLORIDE

? s s1 and s6

40380 S1

942 S6

S7 10 S1 AND S6

? t s7/9,k/all

7/9,K/1 (Item 1 from file: 5)

DIALOG(R)File 5:Biosis Previews(R)

(c) 2007 The Thomson Corporation. All rts. reserv.

0015321550 BIOSIS NO.: 200510016050

Arginine hydrochloride enhances the dynamics of subunit assembly and  
the chaperone-like activity of alpha- crystallin

AUTHOR: Srinivas V; Raman B; Rao K Sridhar; Ramakrishna T; Rao Ch Mohan  
(Reprint)

AUTHOR ADDRESS: Ctr Cellular and Mol Biol, Uppal Rd, Hyderabad 500007,  
Andhra Pradesh, India\*\*India

AUTHOR E-MAIL ADDRESS: mohan@ccmb.res.in  
JOURNAL: Molecular Vision 11 (27-29): p249-255 APR 1 05 2005  
ISSN: 1090-0535  
DOCUMENT TYPE: Article  
RECORD TYPE: Abstract  
LANGUAGE: English

**ABSTRACT:** Purpose: alpha- **Crystallin** , a major eye lens protein, bears homology with small heat shock proteins (sHsps) and exhibits molecular chaperone-like activity. Structural perturbation by temperature or low concentrations of denaturants leads to enhancement of its chaperone-like activity. We have earlier demonstrated similar enhancement of chaperone-like activity using biologically compatible solutes such as **arginine hydrochloride** and aminoguanidine. The purpose of the present study is to get an insight into the mechanism of the arginine induced enhancement of chaperone-like activity of alpha- **crystallin** .Methods: The effect of **arginine hydrochloride** on the chaperone-like activity of alpha- **crystallin** at 25 degrees C was studied using DTT induced aggregation of insulin as a model system. Changes in the accessibility of the thiol group near the end of the alpha- **crystallin** domain in the absence and the presence of **arginine hydrochloride** were studied using dithiobisnitrobenzoic acid. Fluorescence resonance energy transfer studies were performed to investigate changes in the dynamics of the subunit assembly. Urea induced denaturation studies of alpha- **crystallin** were carried out to investigate structural destabilization of alpha- **crystallin** , if any, in the presence of **arginine hydrochloride** .Results: **Arginine hydrochloride** increases the chaperone-like activity of alpha- **crystallin** several fold towards DTT induced aggregation of insulin at room temperature. Our study shows that both the extent and the rate of accessibility of the thiol group are increased in the presence of arginine. Fluorescence resonance energy transfer experiments show that **arginine hydrochloride** significantly increases the subunit exchange between the oligomers of alpha- **crystallin** . Arginine induced structural perturbation and loosening of subunit assembly of alpha- **crystallin** leads to overall destabilization of the protein as reflected by the urea denaturation study.Conclusions: Arginine perturbs the tertiary and quaternary structure of alpha- **crystallin** and enhances the dynamics of the subunit assembly leading to enhanced chaperone-like activity. Thus, in addition to size, surface hydrophobicity, and charge distribution, the dynamics of the subunit assembly appears to be one of the critical factors that can modulate the chaperone activity.

REGISTRY NUMBERS: 9004-10-8: insulin; 79-17-4: aminoguanidine; 32042-43-6: **arginine hydrochloride**

**DESCRIPTORS:**

MAJOR CONCEPTS: Biochemistry and Molecular Biophysics; Sense Organs-- Sensory Reception

ORGANISMS: PARTS ETC: eye lens

CHEMICALS & BIOCHEMICALS: insulin; small heat shock proteins; aminoguanidine; **arginine hydrochloride** ; alpha- **crystallin** -- chaperone-like activity

**CONCEPT CODES:**

10060 Biochemistry studies - General

10064 Biochemistry studies - Proteins, peptides and amino acids

20004 Sense organs - Physiology and biochemistry

**Arginine hydrochloride enhances the dynamics of subunit assembly and the chaperone-like activity of alpha-crystallin**

**ABSTRACT:** Purpose: alpha- Crystallin , a major eye lens protein, bears homology with small heat shock proteins (sHsps) and exhibits...

...have earlier demonstrated similar enhancement of chaperone-like activity using biologically compatible solutes such as arginine hydrochloride and aminoguanidine. The purpose of the present study is to get an insight into the mechanism of the arginine induced enhancement of chaperone-like activity of alpha- crystallin .Methods: The effect of arginine hydrochloride on the chaperone-like activity of alpha- crystallin at 25 degrees C was studied using DTT induced aggregation of insulin as a model system. Changes in the accessibility of the thiol group near the end of the alpha- crystallin domain in the absence and the presence of arginine hydrochloride were studied using dithiobisnitrobenzoic acid. Fluorescence resonance energy transfer studies were performed to investigate changes in the dynamics of the subunit assembly. Urea induced denaturation studies of alpha- crystallin were carried out to investigate structural destabilization of alpha- crystallin , if any, in the presence of arginine hydrochloride .Results: Arginine hydrochloride increases the chaperone-like activity of alpha- crystallin several fold towards DTT induced aggregation of insulin at room temperature. Our study shows that...

...group are increased in the presence of arginine. Fluorescence resonance energy transfer experiments show that arginine hydrochloride significantly increases the subunit exchange between the oligomers of alpha- crystallin . Arginine induced structural perturbation and loosening of subunit assembly of alpha- crystallin leads to overall destabilization of the protein as reflected by the urea denaturation study.Conclusions: Arginine perturbs the tertiary and quaternary structure of alpha- crystallin and enhances the dynamics of the subunit assembly leading to enhanced chaperone-like activity. Thus...

...REGISTRY NUMBERS: arginine hydrochloride

DESCRIPTORS:

CHEMICALS & BIOCHEMICALS: ... arginine hydrochloride ; ...

...alpha- crystallin --

7/9,K/2 (Item 2 from file: 5)

DIALOG(R)File 5:Biosis Previews(R)

(c) 2007 The Thomson Corporation. All rts. reserv.

0014361712 BIOSIS NO.: 200300320431

**Structural perturbation and enhancement of the chaperone-like activity of alpha- crystallin by arginine hydrochloride .**

**AUTHOR:** Srinivas Volety; Raman Bakthisaran; Rao Kunchala Sridhar; Ramakrishna Tangirala; Rao Ch Mohan (Reprint)

**AUTHOR ADDRESS:** Centre for Cellular and Molecular Biology, Uppal Road, Hyderabad, 500 007, India\*\*India

**AUTHOR E-MAIL ADDRESS:** mohan@ccmb.res.in

**JOURNAL:** Protein Science 12 (6): p1262-1270 June 2003 2003

**MEDIUM:** print

**ISSN:** 0961-8368

**DOCUMENT TYPE:** Article

RECORD TYPE: Abstract  
LANGUAGE: English

ABSTRACT: Structural perturbation of alpha- **crystallin** is shown to enhance its molecular chaperone-like activity in preventing aggregation of target proteins. We demonstrate that arginine, a biologically compatible molecule that is known to bind to the peptide backbone and negatively charged side-chains, increases the chaperone-like activity of calf eye lens alpha- **crystallin** as well as recombinant human alphaA- and alphaB-crystallins. Arginine-induced increase in the chaperone activity is more pronounced for alphaB- **crystallin** than for alphaA- **crystallin** . Other guanidinium compounds such as aminoguanidine hydrochloride and guanidine hydrochloride also show a similar effect, but to different extents. A point mutation, R120G, in alphaB- **crystallin** that is associated with desmin-related myopathy, results in a significant loss of chaperone-like activity. Arginine restores the activity of mutant protein to a considerable extent. We have investigated the effect of arginine on the structural changes of alpha- **crystallin** by circular dichroism, fluorescence, and glycerol gradient sedimentation. Far-UV CD spectra show no significant changes in secondary structure, whereas near-UV CD spectra show subtle changes in the presence of arginine. Glycerol gradient sedimentation shows a significant decrease in the size of alpha- **crystallin** oligomer in the presence of arginine. Increased exposure of hydrophobic surfaces of alpha- **crystallin** , as monitored by pyrene-solubilization and ANS-fluorescence, is observed in the presence of arginine. These results show that arginine brings about subtle changes in the tertiary structure and significant changes in the quaternary structure of alpha- **crystallin** and enhances its chaperone-like activity significantly. This study should prove useful in designing strategies to improve chaperone function for therapeutic applications.

REGISTRY NUMBERS: 1119-34-2Q: **arginine hydrochloride** ; 15595-35-4Q: **arginine hydrochloride** ; 32042-43-6Q: **arginine hydrochloride** ; 1937-19-5Q: aminoguanidine hydrochloride; 16139-18-7Q: aminoguanidine hydrochloride

DESCRIPTORS:

MAJOR CONCEPTS: Biochemistry and Molecular Biophysics

BIOSYSTEMATIC NAMES: Bovidae--Artiodactyla, Mammalia, Vertebrata, Chordata, Animalia; Hominidae--Primates, Mammalia, Vertebrata, Chordata, Animalia

ORGANISMS: cow (Bovidae); human (Hominidae)

ORGANISMS: PARTS ETC: lens--sensory system

COMMON TAXONOMIC TERMS: Artiodactyls; Nonhuman Vertebrates; Nonhuman Mammals; Animals; Chordates; Humans; Mammals; Primates; Vertebrates

DISEASES: desmin-related myopathy--muscle disease

CHEMICALS & BIOCHEMICALS: alpha- **crystallin** --structure, chaperone-like activity; **arginine hydrochloride** ; alpha-A **crystallin** ; alpha-B **crystallin** ; aminoguanidine hydrochloride

METHODS & EQUIPMENT: far-UV circular dichroism spectroscopy--laboratory techniques, spectrum analysis techniques; near-UV circular dichroism spectroscopy--laboratory techniques, spectrum analysis techniques; fluorescence assay--laboratory techniques, spectrum analysis techniques; glycerol gradient sedimentation--laboratory techniques

MISCELLANEOUS TERMS: drug development

CONCEPT CODES:

10060 Biochemistry studies - General

17506 Muscle - Pathology

20004 Sense organs - Physiology and biochemistry

BIOSYSTEMATIC CODES:

85715 Bovidae

86215 Hominidae

**Structural perturbation and enhancement of the chaperone-like activity of alpha- crystallin by arginine hydrochloride .**

**ABSTRACT:** Structural perturbation of alpha- crystallin is shown to enhance its molecular chaperone-like activity in preventing aggregation of target proteins...

...and negatively charged side-chains, increases the chaperone-like activity of calf eye lens alpha- crystallin as well as recombinant human alphaA- and alphaB-crystallins. Arginine-induced increase in the chaperone activity is more pronounced for alphaB- crystallin than for alphaA- crystallin . Other guanidinium compounds such as aminoguanidine hydrochloride and guanidine hydrochloride also show a similar effect, but to different extents. A point mutation, R120G, in alphaB- crystallin that is associated with desmin-related myopathy, results in a significant loss of chaperone-like...

...considerable extent. We have investigated the effect of arginine on the structural changes of alpha- crystallin by circular dichroism, fluorescence, and glycerol gradient sedimentation. Far-UV CD spectra show no significant...

...presence of arginine. Glycerol gradient sedimentation shows a significant decrease in the size of alpha- crystallin oligomer in the presence of arginine. Increased exposure of hydrophobic surfaces of alpha- crystallin , as monitored by pyrene-solubilization and ANS-fluorescence, is observed in the presence of arginine...

...subtle changes in the tertiary structure and significant changes in the quaternary structure of alpha- crystallin and enhances its chaperone-like activity significantly. This study should prove useful in designing strategies...

...REGISTRY NUMBERS: arginine hydrochloride ; ...

... arginine hydrochloride ; ...

... arginine hydrochloride ;

DESCRIPTORS:

CHEMICALS & BIOCHEMICALS: alpha- crystallin --...

... arginine hydrochloride ; ...

...alpha-A crystallin ; ...

...alpha-B crystallin ;

7/9,K/3 (Item 1 from file: 34)

DIALOG(R) File 34:SciSearch(R) Cited Ref Sci

(c) 2007 The Thomson Corp. All rts. reserv.

14246175 Genuine Article#: 953HG Number of References: 44

Title: Modulation of alpha- crystallin chaperone activity in diabetic rat

# **lens by curcumin**

Author(s): Kumar PA; Suryanarayana P; Reddy PY; Reddy GB (REPRINT)  
Corporate Source: Natl Inst Nutr, Hyderabad 500007/Andhra Pradesh/India/  
(REPRINT); Natl Inst Nutr, Hyderabad 500007/Andhra Pradesh/India/(  
geereddy@yahoo.com)

Journal: MOLECULAR VISION, 2005, V11, N66 (JUL 26), P561-568

ISSN: 1090-0535 Publication date: 20050726

Publisher: MOLECULAR VISION, C/O JEFF BOATRIGHT, LAB B, 5500 EMORY EYE  
CENTER, 1327 CLIFTON RD, N E, ATLANTA, GA 30322 USA

Language: English Document Type: ARTICLE

Geographic Location: India

Journal Subject Category: BIOCHEMISTRY & MOLECULAR BIOLOGY; OPHTHALMOLOGY

Abstract: Purpose: A decline in the chaperone-like activity of eye lens  
alpha- **crystallin** in diabetic conditions has been reported. In this  
study, we investigated whether curcumin, a dietary antioxidant, can  
manipulate the chaperone-like activity of alpha- **crystallin** in  
diabetic rat lens.

Methods: A group of rats received ip injection of streptozotocin  
(STZ; 35 mg/kg body weight in buffer) to induce hyperglycemia, while  
another group of rats received only buffer as vehicle and served as  
control. STZ-treated rats were assigned to 3 groups and fed either no  
curcumin or 0.002% or 0.01% curcumin, respectively. Cataract  
progression due to hyperglycemia was monitored with a slit lamp  
biomicroscope. At the end of 8 weeks animals were sacrificed and lenses  
were collected. alpha H- and alpha L-crystallins from a set of pooled  
lenses in each group were isolated by gel filtration. Chaperone  
activity, hydrophobicity, and secondary and tertiary structure of alpha  
H- and alpha L-crystallins were assessed by light  
scattering/spectroscopic methods.

Results: A decrease in chaperone-like activity of alpha H- and  
alpha L-crystallins was observed in STZ-treated diabetic rats. The  
declined chaperone-like activity due to hyperglycemia was associated  
with reduced hydrophobicity and altered secondary and tertiary  
structure of alpha H- and alpha L-crystallins. Interestingly, alpha H-  
and alpha L-crystallins isolated from curcumin fed diabetic rat lenses  
had shown improved chaperone-like activity as compared to alpha H- and  
alpha L-crystallins from untreated diabetic rat lens. Feeding of  
curcumin prevented the alterations in hydrophobicity and structural  
changes due to STZ-induced hyperglycemia. Modulation of functional and  
structural properties by curcumin was found to be greater with the  
alpha L- **crystallin** than alpha H- **crystallin** . Loss of chaperone  
activity of alpha- **crystallin** , particularly alpha L- **crystallin** , in  
diabetic rat lens could be attributed at least partly to increased  
oxidative stress. Being an antioxidant, curcumin feeding has prevented  
the loss of alpha- **crystallin** chaperone activity and delayed the  
progression and maturation of diabetic cataract.

Conclusions: We demonstrate that curcumin, at the levels close to  
dietary consumption, prevented the loss of chaperone-like activity of  
alpha- **crystallin** vis-a-vis cataractogenesis due to diabetes in rat  
lens.

Identifiers--KeyWord Plus(R): A- **CRYSTALLIN** ; B- **CRYSTALLIN** ; **ARGININE**  
**HYDROCHLORIDE** ; IN-VIVO; CATARACT; AGGREGATION; PROTECT; STRESS; INDIA;  
RISK

Cited References:

- \*INT DIAB FED, 2001, DIAB ATL 2000  
BAI F, 2003, V278, P36876, J BIOL CHEM  
BHAT SP, 2003, V60, P205, PROG DRUG RES  
BOYLE DL, 2003, V3, P3, BMC OPHTHALMOL  
BRADY JP, 1997, V94, P884, P NATL ACAD SCI USA  
CHERIAN M, 1995, V212, P184, BIOCHEM BIOPH RES CO  
CHERIAN M, 1995, V208, P675, BIOCHEM BIOPH RES CO  
CLARK JI, 1996, V93, P15185, P NATL ACAD SCI USA  
DAS KP, 1995, V369, P321, FEBS LETT  
DERHAM BK, 2002, V364, P711, BIOCHEM J 3  
DERHAM BK, 1997, V328, P763, BIOCHEM J 3  
GANEVA E, 2001, V2, P205, CURR PROTEIN PEPT SC  
GROENEN PJTA, 1994, V225, P1, EUR J BIOCHEM  
HANSON SRA, 2000, V71, P195, EXP EYE RES  
HARDING J, 1991, CATARACT BIOICM EPID  
HOOK DWA, 1997, V247, P380, EUR J BIOCHEM  
HORWITZ J, 1998, V290, P365, METHOD ENZYMOL  
HORWITZ J, 1992, V11, P817, CURR EYE RES  
HORWITZ J, 2003, V76, P145, EXP EYE RES  
HUANG FY, 2000, V273, P197, BIOCHEM BIOPH RES CO  
KADOR PF, 1985, V28, P841, J MED CHEM  
KUMAR MS, 2005, V280, P21726, J BIOL CHEM  
KUMAR MS, 2004, V379, P273, BIOCHEM J 2  
KYSELOVA Z, 2004, V18, P129, J DIABETES COMPLICAT  
MARES JA, 2004, V62, P28, NUTR REV  
NIRMALAN PK, 2004, V88, P989, BRIT J OPHTHALMOL  
PRADEEPA R, 2002, V116, P121, INDIAN J MED RES  
RAMAN B, 1997, V272, P23559, J BIOL CHEM  
REDDY GB, 2000, V275, P4565, J BIOL CHEM  
REDDY G, 2002, V8, P298, MOL VIS  
REDDY GB, 2001, V282, P712, BIOCHEM BIOPH RES CO  
REEVES PG, 1997, V127, P838, J NUTR  
SHARMA KK, 1998, V273, P8965, J BIOL CHEM  
SPECTOR A, 2000, V16, P193, J OCUL PHARMACOL TH  
SRINIVAS V, 2003, V12, P1262, PROTEIN SCI  
SRINIVAS V, 2005, V11, P249, MOL VIS  
SURYANARAYANA P, 2003, V9, P223, MOL VIS  
SURYANARAYANA P, 2005, V46, P2092, INVEST OPHTH VIS SCI  
THAMPI P, 2002, V229, P113, MOL CELL BIOCHEM  
THAMPI P, 2002, V43, P3265, INVEST OPHTH VIS SCI  
THAMPI P, 2003, V42, P11857, BIOCHEMISTRY-US  
UGHADE SN, 1998, V46, P221, INDIAN J OPHTHALMOL  
VANBOEKEL MAM, 1992, V314, P1, FEBS LETT  
YANG FS, 2005, V280, P5892, J BIOL CHEM

**Title: Modulation of alpha- crystallin chaperone activity in diabetic rat lens by curcumin**

**Abstract:** Purpose: A decline in the chaperone-like activity of eye lens alpha- crystallin in diabetic conditions has been reported. In this study, we investigated whether curcumin, a dietary antioxidant, can manipulate the chaperone-like activity of alpha- crystallin in diabetic rat lens.

Methods: A group of rats received ip injection of streptozotocin (STZ...

...functional and structural properties by curcumin was found to be greater with the alpha L- **crystallin** than alpha H- **crystallin** . Loss of chaperone activity of alpha- **crystallin** , particularly alpha L- **crystallin** , in diabetic rat lens could be attributed at least partly to increased oxidative stress. Being an antioxidant, curcumin feeding has prevented the loss of alpha- **crystallin** chaperone activity and delayed the progression and maturation of diabetic cataract.

Conclusions: We demonstrate that...

...the levels close to dietary consumption, prevented the loss of chaperone-like activity of alpha- **crystallin** vis-a-vis cataractogenesis due to diabetes in rat lens.

...Identifiers--A- **CRYSTALLIN** ; B- **CRYSTALLIN** ; **ARGININE HYDROCHLORIDE** ; IN-VIVO; CATARACT; AGGREGATION; PROTECT; STRESS; INDIA; RISK

7/9,K/4 (Item 2 from file: 34)

DIALOG(R) File 34:SciSearch(R) Cited Ref Sci  
(c) 2007 The Thomson Corp. All rts. reserv.

13835968 Genuine Article#: 915KO Number of References: 49

**Title:** Arginine hydrochloride enhances the dynamics of subunit assembly and the chaperone-like activity of alpha- crystallin

**Author(s):** Srinivas V; Raman B; Rao KS; Ramakrishna T; Rao CM (REPRINT)

**Corporate Source:** Ctr Cellular & Mol Biol,Uppal Rd/Hyderabad 500007/Andhra Pradesh/India/ (REPRINT); Ctr Cellular & Mol Biol,Hyderabad 500007/Andhra Pradesh/India/(mohan@ccmb.res.in)

**Journal:** MOLECULAR VISION, 2005, V11, N27-29 (APR 1), P249-255

**ISSN:** 1090-0535 **Publication date:** 20050401

**Publisher:** MOLECULAR VISION, C/O JEFF BOATRIGHT, LAB B, 5500 EMORY EYE CENTER, 1327 CLIFTON RD, N E, ATLANTA, GA 30322 USA

**Language:** English **Document Type:** ARTICLE

**Geographic Location:** India

**Journal Subject Category:** BIOCHEMISTRY & MOLECULAR BIOLOGY; OPHTHALMOLOGY

**Abstract:** Purpose: alpha- **Crystallin** , a major eye lens protein, bears homology with small heat shock proteins (sHsps) and exhibits molecular chaperone-like activity. Structural perturbation by temperature or low concentrations of denaturants leads to enhancement of its chaperone-like activity. We have earlier demonstrated similar enhancement of chaperone-like activity using biologically compatible solutes such as **arginine hydrochloride** and aminoguanidine. The purpose of the present study is to get an insight into the mechanism of the arginine induced enhancement of chaperone-like activity of alpha- **crystallin** .

**Methods:** The effect of **arginine hydrochloride** on the chaperone-like activity of alpha- **crystallin** at 25 degrees C was studied using DTT induced aggregation of insulin as a model system. Changes in the accessibility of the thiol group near the end of the alpha- **crystallin** domain in the absence and the presence of **arginine hydrochloride** were studied using dithiobisnitrobenzoic acid. Fluorescence resonance energy transfer studies were performed to investigate changes in the dynamics of the subunit assembly. Urea induced denaturation studies of alpha- **crystallin** were carried out to investigate structural destabilization of alpha- **crystallin** , if any, in the presence of **arginine hydrochloride** .



Results: **Arginine hydrochloride** increases the chaperone-like activity of **alpha-crystallin** several fold towards DTT induced aggregation of insulin at room temperature. Our study shows that both the extent and the rate of accessibility of the thiol group are increased in the presence of arginine. Fluorescence resonance energy transfer experiments show that **arginine hydrochloride** significantly increases the subunit exchange between the oligomers of **alpha-crystallin**. Arginine induced structural perturbation and loosening of subunit assembly of **alpha-crystallin** leads to overall destabilization of the protein as reflected by the urea denaturation study.

Conclusions: Arginine perturbs the tertiary and quaternary structure of **alpha-crystallin** and enhances the dynamics of the subunit assembly leading to enhanced chaperone-like activity. Thus, in addition to size, surface hydrophobicity, and charge distribution, the dynamics of the subunit assembly appears to be one of the critical factors that can modulate the chaperone activity.

Identifiers--KeyWord Plus(R): HEAT-SHOCK-PROTEIN; DESMIN-RELATED MYOPATHY; QUATERNARY STRUCTURE; MOLECULAR CHAPERONE; A- **CRYSTALLIN** ; B- **CRYSTALLIN** ; STRUCTURAL PERTURBATION; HYDROPHOBIC SURFACES; MISSENSE MUTATION; ENERGY-TRANSFER

Cited References:

- AUGUSTEYN RC, 1987, V915, P132, BIOCHIM BIOPHYS ACTA  
BENNDORF R, 2001, V276, P26753, J BIOL CHEM  
BOVA MP, 2002, V277, P38468, J BIOL CHEM  
BOVA MP, 1999, V96, P6137, P NATL ACAD SCI USA  
BOVA MP, 2000, V275, P1035, J BIOL CHEM  
BOVA MP, 1997, V272, P29511, J BIOL CHEM  
DAS BK, 1997, V236, P370, BIOCHEM BIOPH RES CO  
DAS KP, 1995, V369, P321, FEBS LETT  
DAS KP, 1995, V311, P367, BIOCHEM J 2  
DATTA SA, 2000, V275, P41004, J BIOL CHEM  
DEJONG WW, 1998, V22, P151, INT J BIOL MACROMOL  
GANEA E, 2000, V345, P467, BIOCHEM J 3  
GOENKA S, 2001, V359, P547, BIOCHEM J 3  
GU LX, 2002, V319, P517, J MOL BIOL  
HASLBECK M, 1999, V18, P6744, EMBO J  
HESS JF, 1998, V4, P29, MOL VIS  
HIRATSUKA T, 1992, V267, P14941, J BIOL CHEM  
HOOK DWA, 1997, V247, P380, EUR J BIOCHEM  
HORWITZ J, 1992, V89, P10449, P NATL ACAD SCI USA  
INGOLIA TD, 1982, V79, P2360, P NATL ACAD SCI-BIOL  
KLEMENZ R, 1991, V88, P3652, P NATL ACAD SCI USA  
KUMAR LVS, 1999, V274, P24137, J BIOL CHEM  
LAMBERT H, 1999, V274, P9378, J BIOL CHEM  
LEUNG SM, 1996, V1, P78, CELL STRESS CHAPERON  
LITT M, 1998, V7, P471, HUM MOL GENET  
MARINI I, 2000, V275, P32559, J BIOL CHEM  
NARBERHAUS F, 2002, V66, P64, MICROBIOL MOL BIOL R  
PERNG MD, 1999, V274, P33235, J BIOL CHEM  
RAJARAMAN K, 2001, V497, P118, FEBS LETT  
RAMAN B, 1997, V272, P23559, J BIOL CHEM  
RAMAN B, 1994, V269, P27264, J BIOL CHEM  
RAO CM, 2002, V68, P349, P IND NATL SCI ACAD  
RAO CM, 1998, V22, P271, INT J BIOL MACROMOL  
RAWAT U, 1998, V273, P9415, J BIOL CHEM

RIDDLES PW, 1983, V91, P49, METHOD ENZYMOL  
 SIEZEN RJ, 1978, V537, P456, BIOCHIM BIOPHYS ACTA  
 SIEZEN RJ, 1980, V111, P435, EUR J BIOCHEM  
 SIEZEN RJ, 1978, V91, P387, EUR J BIOCHEM  
 SMITH JB, 1996, V63, P125, EXP EYE RES  
 SRINIVAS V, 2003, V12, P1262, PROTEIN SCI  
 SUN TX, 1998, V430, P401, FEBS LETT  
 VANDENOETELAAR PJM, 1990, V29, P3488, BIOCHEMISTRY-US  
 VANHOUDT J, 2000, V39, P4483, BIOCHEMISTRY-US  
 VANMONTFORT RLM, 2001, V8, P1025, NAT STRUCT BIOL  
 VANMONTFORT R, 2002, V59, P105, ADV PROTEIN CHEM  
 VICART P, 1998, V20, P92, NAT GENET  
 WANG KY, 1994, V269, P13601, J BIOL CHEM  
 YANG HM, 1999, V8, P174, PROTEIN SCI  
 YONEHARA M, 1996, V271, P2641, J BIOL CHEM

**Title:** Arginine hydrochloride enhances the dynamics of subunit assembly and the chaperone-like activity of alpha-crystallin.

**Abstract:** Purpose: alpha-Crystallin, a major eye lens protein, bears homology with small heat shock proteins (sHsps) and exhibits...

...have earlier demonstrated similar enhancement of chaperone-like activity using biologically compatible solutes such as arginine hydrochloride and aminoguanidine. The purpose of the present study is to get an insight into the mechanism of the arginine induced enhancement of chaperone-like activity of alpha-crystallin.

**Methods:** The effect of arginine hydrochloride on the chaperone-like activity of alpha-crystallin at 25 degrees C was studied using DTT induced aggregation of insulin as a model system. Changes in the accessibility of the thiol group near the end of the alpha-crystallin domain in the absence and the presence of arginine hydrochloride were studied using dithiobisnitrobenzoic acid. Fluorescence resonance energy transfer studies were performed to investigate changes in the dynamics of the subunit assembly. Urea induced denaturation studies of alpha-crystallin were carried out to investigate structural destabilization of alpha-crystallin, if any, in the presence of arginine hydrochloride.

**Results:** Arginine hydrochloride increases the chaperone-like activity of alpha-crystallin several fold towards DTT induced aggregation of insulin at room temperature. Our study shows that...

...group are increased in the presence of arginine. Fluorescence resonance energy transfer experiments show that arginine hydrochloride significantly increases the subunit exchange between the oligomers of alpha-crystallin. Arginine induced structural perturbation and loosening of subunit assembly of alpha-crystallin leads to overall destabilization of the protein as reflected by the urea denaturation study.

**Conclusions:** Arginine perturbs the tertiary and quaternary structure of alpha-crystallin and enhances the dynamics of the subunit assembly leading to enhanced chaperone-like activity. Thus...

...Identifiers--HEAT-SHOCK-PROTEIN; DESMIN-RELATED MYOPATHY; QUATERNARY STRUCTURE; MOLECULAR CHAPERONE; A-CRYSTALLIN; B-CRYSTALLIN; STRUCTURAL PERTURBATION; HYDROPHOBIC SURFACES; MISSENSE MUTATION; ENERGY-TRANSFER

7/9,K/5 (Item 3 from file: 34)

DIALOG(R)File 34:SciSearch(R) Cited Ref Sci  
(c) 2007 The Thomson Corp. All rts. reserv.

11679571 Genuine Article#: 681LC Number of References: 57

**Title: Structural perturbation and enhancement of the chaperone-like activity of alpha- crystallin by arginine hydrochloride**

Author(s): Srinivas V; Raman B; Rao KS; Ramakrishna T; Rao CM (REPRINT)

Corporate Source: Ctr Cellular & Mol Biol,Uppal Rd/Hyderabad 500007/Andhra Pradesh/India/ (REPRINT); Ctr Cellular & Mol Biol,Hyderabad 500007/Andhra Pradesh/India/

Journal: PROTEIN SCIENCE, 2003, V12, N6 (JUN), P1262-1270

ISSN: 0961-8368 Publication date: 20030600

Publisher: COLD SPRING HARBOR LAB PRESS, PUBLICATIONS DEPT, 500 SUNNYSIDE BLVD, WOODBURY, NY 11797-2924 USA

Language: English Document Type: ARTICLE

Geographic Location: India

Journal Subject Category: BIOCHEMISTRY & MOLECULAR BIOLOGY

**Abstract:** Structural perturbation of  $\alpha$ - **crystallin** is shown to enhance its molecular chaperone-like activity in preventing aggregation of target proteins. We demonstrate that arginine, a biologically compatible molecule that is known to bind to the peptide backbone and negatively charged side-chains, increases the chaperone-like activity of calf eye lens  $\alpha$ - **crystallin** as well as recombinant human  $\alpha$ A- and  $\alpha$ B-**crystallins**. Arginine-induced increase in the chaperone activity is more pronounced for  $\alpha$ B- **crystallin** than for  $\alpha$ A- **crystallin**. Other guanidinium compounds such as aminoguanidine hydrochloride and guanidine hydrochloride also show a similar effect, but to different extents. A point mutation, R120G, in  $\alpha$ B- **crystallin** that is associated with desmin-related myopathy, results in a significant loss of chaperone-like activity. Arginine restores the activity of mutant protein to a considerable extent. We have investigated the effect of arginine on the structural changes of  $\alpha$ - **crystallin** by circular dichroism, fluorescence, and glycerol gradient sedimentation. Far-UV CD spectra show no significant changes in secondary structure, whereas near-UV CD spectra show subtle changes in the presence of arginine. Glycerol gradient sedimentation shows a significant decrease in the size of  $\alpha$ - **crystallin** oligomer in the presence of arginine. Increased exposure of hydrophobic surfaces of  $\alpha$ - **crystallin**, as monitored by pyrene-solubilization and ANS-fluorescence, is observed in the presence of arginine. These results show that arginine brings about subtle changes in the tertiary structure and significant changes in the quaternary structure of  $\alpha$ - **crystallin** and enhances its chaperone-like activity significantly. This study should prove useful in designing strategies to improve chaperone function for therapeutic applications.

**Descriptors--Author Keywords:** chaperone-like activity ;  $\alpha$ - **crystallin** ; arginine ; aminoguanidine ; structural perturbation

**Identifiers--KeyWord Plus(R):** HEAT-SHOCK-PROTEIN; DESMIN-RELATED MYOPATHY; B- **CRYSTALLIN** ; MOLECULAR CHAPERONE; A- **CRYSTALLIN** ; IN-VITRO; HYDROPHOBIC SURFACES; MISSENSE MUTATION; THERMAL-STRESS; LENS

**Cited References:**

AOYAMA A, 1993, V55, P760, INT J CANCER

ARAKAWA T, 1984, V23, P5924, BIOCHEMISTRY-US

BHAT SP, 1989, V158, P319, BIOCHEM BIOPH RES CO

BORKMAN RF, 1996, V62, P141, EXP EYE RES

BOVA MP, 1999, V96, P6137, P NATL ACAD SCI USA  
 CLARK JI, 1996, V93, P15185, P NATL ACAD SCI USA  
 CRAIG EA, 1991, V16, P135, TRENDS BIOCHEM SCI  
 DAS KP, 1995, V369, P321, FEBS LETT  
 DAS BK, 1997, V236, P370, BIOCHEM BIOPH RES CO  
 DATTA SA, 1999, V274, P34773, J BIOL CHEM  
 DIAMANT S, 2001, V276, P39586, J BIOL CHEM  
 DUBIN RA, 1989, V9, P1083, MOL CELL BIOL  
 GANEA E, 2000, V345, P467, BIOCHEM J 3  
 GOENKA S, 2001, V359, P547, BIOCHEM J 3  
 GROENEN PJTA, 1994, V225, P1, EUR J BIOCHEM  
 GU LX, 2002, V319, P517, J MOL BIOL  
 HAKLAR G, 1998, V25, P908, CLIN EXP PHARMACOL P  
 HASLBECK M, 1999, V18, P6744, EMBO J  
 HESS JF, 1998, V4, P29, MOL VIS  
 HOOK DWA, 1997, V247, P380, EUR J BIOCHEM  
 HORWITZ J, 1992, V89, P10449, P NATL ACAD SCI USA  
 INGOLIA TD, 1982, V79, P2360, P NATL ACAD SCI USA  
 INOMATA M, 2000, V128, P771, J BIOCHEM-TOKYO  
 IWAKI T, 1989, V57, P71, CELL  
 KATO K, 1991, V1080, P173, BIOCHIM BIOPHYS ACTA  
 KUMAR LVS, 1999, V274, P24137, J BIOL CHEM  
 LAMBERT H, 1999, V274, P9378, J BIOL CHEM  
 LEUNG SM, 1996, V1, P78, CELL STRESS CHAPERON  
 LIN TY, 1996, V5, P372, PROTEIN SCI  
 LITT M, 1998, V7, P471, HUM MOL GENET  
 MARINI I, 2000, V275, P32559, J BIOL CHEM  
 MERCK KB, 1993, V268, P1046, J BIOL CHEM  
 PALMISANO DV, 1995, V1246, P91, BBA-PROTEIN STRUCT M  
 RAJARAMAN K, 2001, V497, P118, FEBS LETT  
 RAMAN B, 1994, V269, P27264, J BIOL CHEM  
 RAO PV, 1992, V54, P627, EXP EYE RES  
 RAO CM, 1998, V22, P271, INT J BIOL MACROMOL  
 RAO CM, 2002, V68, P349, P IND NATL SCI ACAD  
 RAWAT U, 1998, V273, P9415, J BIOL CHEM  
 REDDY GB, 2000, V275, P4565, J BIOL CHEM  
 RENKAWEK K, 1994, V87, P155, ACTA NEUROPATHOL  
 RUDOLPH R, 1996, V10, P49, FASEB J  
 SHARMA KK, 2000, V275, P3767, J BIOL CHEM  
 SHARMA KK, 1997, V239, P217, BIOCHEM BIOPH RES CO  
 SMITH JB, 1996, V63, P125, EXP EYE RES  
 SONG JL, 2001, V276, P40241, J BIOL CHEM  
 STRYER L, 1965, V13, P482, J MOL BIOL  
 SUEMATSU Y, 2001, V19, P873, EUR J CARDIO-THORAC  
 SULOCHANA KN, 1998, V67, P597, EXP EYE RES  
 SUN TX, 1997, V272, P6220, J BIOL CHEM  
 SWAMYMRUTHINTI S, 1996, V62, P505, EXP EYE RES  
 TIEMAN BC, 2001, V276, P44541, J BIOL CHEM  
 TIMASHEFF SN, 1988, P331, PROTEIN STRUCTURE PR  
 VANMONTFORT RLM, 2001, V8, P1025, NAT STRUCT BIOL  
 VICART P, 1998, V20, P92, NAT GENET  
 VOZIYAN PA, 2000, V89, P1036, J PHARM SCI  
 YANG HM, 1999, V8, P174, PROTEIN SCI

**Title: Structural perturbation and enhancement of the chaperone-like activity of alpha- crystallin by arginine hydrochloride**

**Abstract: Structural perturbation of a- crystallin is shown to enhance its**

molecular chaperone-like activity in preventing aggregation of target proteins...

- ...and negatively charged side-chains, increases the chaperone-like activity of calf eye lens alpha- **crystallin** as well as recombinant human alphaA- and alphaB-crystallins. Arginine-induced increase in the chaperone activity is more pronounced for alphaB- **crystallin** than for alphaA- **crystallin** . Other guanidinium compounds such as aminoguanidine hydrochloride and guanidine hydrochloride also show a similar effect, but to different extents. A point mutation, R120G, in alphaB- **crystallin** that is associated with desmin-related myopathy, results in a significant loss of chaperone-like...
- ...considerable extent. We have investigated the effect of arginine on the structural changes of alpha- **crystallin** by circular dichroism, fluorescence, and glycerol gradient sedimentation. Far-UV CD spectra show no significant...
- ...presence of arginine. Glycerol gradient sedimentation shows a significant decrease in the size of alpha- **crystallin** oligomer in the presence of arginine. Increased exposure of hydrophobic surfaces of alpha- **crystallin** , as monitored by pyrene-solubilization and ANS-fluorescence, is observed in the presence of arginine...
- ...subtle changes in the tertiary structure and significant changes in the quaternary structure of alpha- **crystallin** and enhances its chaperone-like activity significantly. This study should prove useful in designing strategies...
- ...Identifiers--HEAT-SHOCK-PROTEIN; DESMIN-RELATED MYOPATHY; B- **CRYSTALLIN** ; MOLECULAR CHAPERONE; A- **CRYSTALLIN** ; IN-VITRO; HYDROPHOBIC SURFACES; MISSENSE MUTATION; THERMAL-STRESS; LENS

7/9,K/6 (Item 1 from file: 71)

DIALOG(R) File 71:ELSEVIER BIOBASE

(c) 2007 Elsevier B.V. All rts. reserv.

02344455 2003128140

**Structural perturbation and enhancement of the chaperone-like activity of alpha- crystallin by arginine hydrochloride**

Srinivas V.; Raman B.; Rao K.S.; Ramakrishna T.; Rao Ch.M.

ADDRESS: Ch.M. Rao, Ctr. for Cell. and Molecular Biology, Uppal Road, Hyderabad 500 007, India

EMAIL: mohan@ccmb.res.in

Journal: Protein Science, 12/6 (1262-1270), 2003, United States

PUBLICATION DATE: June 1, 2003

CODEN: PRCIE

ISSN: 0961-8368

DOCUMENT TYPE: Article

LANGUAGES: English SUMMARY LANGUAGES: English

NO. OF REFERENCES: 57

Structural perturbation of alpha- **crystallin** is shown to enhance its molecular chaperone-like activity in preventing aggregation of target proteins. We demonstrate that arginine, a biologically compatible molecule that is known to bind to the peptide backbone and negatively charged side-chains, increases the chaperone-like activity of calf eye lens alpha-

**crystallin** as well as recombinant human alphaA- and alphaB-crystallins. Arginine-induced increase in the chaperone activity is more pronounced for alphaB- **crystallin** than for alphaA- **crystallin** . Other guanidinium compounds such as aminoguanidine hydrochloride and guanidine hydrochloride also show a similar effect, but to different extents. A point mutation, R120G, in alphaB- **crystallin** that is associated with desmin-related myopathy, results in a significant loss of chaperone-like activity. Arginine restores the activity of mutant protein to a considerable extent. We have investigated the effect of arginine on the structural changes of alpha- **crystallin** by circular dichroism, fluorescence, and glycerol gradient sedimentation. Far-UV CD spectra show no significant changes in secondary structure, whereas near-UV CD spectra show subtle changes in the presence of arginine. Glycerol gradient sedimentation shows a significant decrease in the size of alpha- **crystallin** oligomer in the presence of arginine. Increased exposure of hydrophobic surfaces of alpha- **crystallin** , as monitored by pyrene-solubilization and ANS-fluorescence, is observed in the presence of arginine. These results show that arginine brings about subtle changes in the tertiary structure and significant changes in the quaternary structure of alpha- **crystallin** and enhances its chaperone-like activity significantly. This study should prove useful in designing strategies to improve chaperone function for therapeutic applications.

DESCRIPTORS:

Chaperone-like activity; alpha- **crystallin** ; Arginine; Aminoguanidine; Structural perturbation

CLASSIFICATION CODE AND DESCRIPTION:

82.2.8 - PROTEIN BIOCHEMISTRY / STRUCTURAL STUDIES / Folding, Unfolding and Stability

**Structural perturbation and enhancement of the chaperone-like activity of alpha- crystallin by arginine hydrochloride**

Structural perturbation of alpha- **crystallin** is shown to enhance its molecular chaperone-like activity in preventing aggregation of target proteins...

...and negatively charged side-chains, increases the chaperone-like activity of calf eye lens alpha- **crystallin** as well as recombinant human alphaA- and alphaB-crystallins. Arginine-induced increase in the chaperone activity is more pronounced for alphaB- **crystallin** than for alphaA- **crystallin** . Other guanidinium compounds such as aminoguanidine hydrochloride and guanidine hydrochloride also show a similar effect, but to different extents. A point mutation, R120G, in alphaB- **crystallin** that is associated with desmin-related myopathy, results in a significant loss of chaperone-like...

...considerable extent. We have investigated the effect of arginine on the structural changes of alpha- **crystallin** by circular dichroism, fluorescence, and glycerol gradient sedimentation. Far-UV CD spectra show no significant...

...presence of arginine. Glycerol gradient sedimentation shows a significant decrease in the size of alpha- **crystallin** oligomer in the presence of arginine. Increased exposure of hydrophobic surfaces of alpha- **crystallin** , as monitored by pyrene-solubilization and ANS-fluorescence, is observed in the presence of arginine...

...subtle changes in the tertiary structure and significant changes in the quaternary structure of alpha- **crystallin** and enhances its chaperone-like activity significantly. This study should prove useful in designing strategies...

DESCRIPTORS:

Chaperone-like activity; alpha- **crystallin** ; Arginine; Aminoguanidine; Structural perturbation

7/9,K/7 (Item 1 from file: 73)

DIALOG(R)File 73:EMBASE

(c) 2007 Elsevier B.V. All rts. reserv.

13357267 EMBASE No: 2005431360

**Arginine hydrochloride enhances the dynamics of subunit assembly and the chaperone-like activity of alpha- crystallin**

Srinivas V.; Raman B.; Sridhar Rao K.; Ramakrishna T.; Mohan Rao Ch.  
Dr. Ch. Mohan Rao, Centre for Cellular and Molecular Biology, Uppal Road,  
Hyderabad 500 007 India

AUTHOR EMAIL: mohan@ccmb.res.in

Molecular Vision ( MOL. VISION ) (United States) 01 APR 2005, 11/-  
(249-255)

CODEN: MVEPF ISSN: 1090-0535

DOCUMENT TYPE: Journal ; Article

LANGUAGE: ENGLISH SUMMARY LANGUAGE: ENGLISH

NUMBER OF REFERENCES: 49

Purpose: alpha- **Crystallin** , a major eye lens protein, bears homology with small heat shock proteins (sHsps) and exhibits molecular chaperone-like activity. Structural perturbation by temperature or low concentrations of denaturants leads to enhancement of its chaperone-like activity. We have earlier demonstrated similar enhancement of chaperone-like activity using biologically compatible solutes such as **arginine hydrochloride** and aminoguanidine. The purpose of the present study is to get an insight into the mechanism of the arginine induced enhancement of chaperone-like activity of **crystallin** . Methods: The effect of **arginine hydrochloride** on the chaperone-like activity of alpha- **crystallin** at 25 degreesC was studied using DTT induced aggregation of insulin as a model system. Changes in the accessibility of the thiol group near the end of the a- **crystallin** domain in the absence and the presence of **arginine hydrochloride** were studied using dithiobisnitrobenzoic acid. Fluorescence resonance energy transfer studies were performed to investigate changes in the dynamics of the subunit assembly. Urea induced denaturation studies of alpha- **crystallin** were carried out to investigate structural destabilization of alpha- **crystallin** , if any, in the presence of **arginine hydrochloride** . Results: **Arginine hydrochloride** increases the chaperone-like activity of alpha- **crystallin** several fold towards DTT induced aggregation of insulin at room temperature. Our study shows that both the extent and the rate of accessibility of the thiol group are increased in the presence of arginine. Fluorescence resonance energy transfer experiments show that **arginine hydrochloride** significantly increases the subunit exchange between the oligomers of alpha- **crystallin** . Arginine induced structural perturbation and loosening of subunit assembly of alpha- **crystallin** leads to overall destabilization of the protein as reflected by the urea denaturation study. Conclusions: Arginine perturbs

the tertiary and quaternary structure of **a-crystallin** and enhances the dynamics of the subunit assembly leading to enhanced chaperone-like activity. Thus, in addition to size, surface hydrophobicity, and charge distribution, the dynamics of the subunit assembly appears to be one of the critical factors that can modulate the chaperone activity. (c)2005 Molecular Vision.

DRUG DESCRIPTORS:

\*arginine; \*chaperone; \*alpha **crystallin**  
dithiothreitol; insulin; benzoic acid; urea

MEDICAL DESCRIPTORS:

protein assembly; biological model; protein domain; fluorescence resonance energy transfer; protein denaturation; protein structure; room temperature; hydrophobicity; article; priority journal

CAS REGISTRY NO.: 1119-34-2, 15595-35-4, 7004-12-8, 74-79-3 (arginine);  
3483-12-3 (dithiothreitol); 9004-10-8 (insulin); 532-32-1, 582-25-2,  
65-85-0, 766-76-7 (benzoic acid); 57-13-6 (urea)

SECTION HEADINGS:

029 Clinical and Experimental Biochemistry

**Arginine hydrochloride enhances the dynamics of subunit assembly and the chaperone-like activity of alpha-crystallin**

Purpose: alpha- **Crystallin** , a major eye lens protein, bears homology with small heat shock proteins (sHsps) and exhibits...

...have earlier demonstrated similar enhancement of chaperone-like activity using biologically compatible solutes such as **arginine hydrochloride** and aminoguanidine. The purpose of the present study is to get an insight into the mechanism of the arginine induced enhancement of chaperone-like activity of **crystallin** . Methods: The effect of **arginine hydrochloride** on the chaperone-like activity of alpha- **crystallin** at 25 degreesC was studied using DTT induced aggregation of insulin as a model system. Changes in the accessibility of the thiol group near the end of the a- **crystallin** domain in the absence and the presence of **arginine hydrochloride** were studied using dithiobisnitrobenzoic acid. Fluorescence resonance energy transfer studies were performed to investigate changes in the dynamics of the subunit assembly. Urea induced denaturation studies of alpha- **crystallin** were carried out to investigate structural destabilization of alpha- **crystallin** , if any, in the presence of **arginine hydrochloride** . Results: **Arginine hydrochloride** increases the chaperone-like activity of alpha- **crystallin** several fold towards DTT induced aggregation of insulin at room temperature. Our study shows that...

...group are increased in the presence of arginine. Fluorescence resonance energy transfer experiments show that **arginine hydrochloride** significantly increases the subunit exchange between the oligomers of alpha- **crystallin** . Arginine induced structural perturbation and loosening of subunit assembly of alpha- **crystallin** leads to overall destabilization of the protein as reflected by the urea denaturation study. Conclusions: Arginine perturbs the tertiary and quaternary structure of a- **crystallin** and enhances the dynamics of the subunit assembly leading to enhanced chaperone-like activity. Thus...

DRUG DESCRIPTORS:

\*arginine; \*chaperone; \*alpha **crystallin**



DIALOG(R) File 73:EMBASE

(c) 2007 Elsevier B.V. All rts. reserv.

12095641 EMBASE No: 2003207170

**Structural perturbation and enhancement of the chaperone-like activity of alpha-crystallin by arginine hydrochloride**

Srinivas V.; Raman B.; Rao K.S.; Ramakrishna T.; Rao Ch.M.

Ch.M. Rao, Ctr. for Cell. and Molecular Biology, Uppal Road, Hyderabad  
500 007 India

AUTHOR EMAIL: mohan@ccmb.res.in

Protein Science ( PROTEIN SCI. ) (United States) 01 JUN 2003, 12/6  
(1262-1270)

CODEN: PRCIE ISSN: 0961-8368

DOCUMENT TYPE: Journal ; Article

LANGUAGE: ENGLISH SUMMARY LANGUAGE: ENGLISH

NUMBER OF REFERENCES: 57

Structural perturbation of alpha-crystallin is shown to enhance its molecular chaperone-like activity in preventing aggregation of target proteins. We demonstrate that arginine, a biologically compatible molecule that is known to bind to the peptide backbone and negatively charged side-chains, increases the chaperone-like activity of calf eye lens alpha-crystallin as well as recombinant human alphaA- and alphaB-crystallins. Arginine-induced increase in the chaperone activity is more pronounced for alphaB-crystallin than for alphaA-crystallin. Other guanidinium compounds such as aminoguanidine hydrochloride and guanidine hydrochloride also show a similar effect, but to different extents. A point mutation, R120G, in alphaB-crystallin that is associated with desmin-related myopathy, results in a significant loss of chaperone-like activity. Arginine restores the activity of mutant protein to a considerable extent. We have investigated the effect of arginine on the structural changes of alpha-crystallin by circular dichroism, fluorescence, and glycerol gradient sedimentation. Far-UV CD spectra show no significant changes in secondary structure, whereas near-UV CD spectra show subtle changes in the presence of arginine. Glycerol gradient sedimentation shows a significant decrease in the size of alpha-crystallin oligomer in the presence of arginine. Increased exposure of hydrophobic surfaces of alpha-crystallin, as monitored by pyrene-solubilization and ANS-fluorescence, is observed in the presence of arginine. These results show that arginine brings about subtle changes in the tertiary structure and significant changes in the quaternary structure of alpha-crystallin and enhances its chaperone-like activity significantly. This study should prove useful in designing strategies to improve chaperone function for therapeutic applications.

**DRUG DESCRIPTORS:**

\*chaperone; \*alpha crystallin --endogenous compound--ec; \*arginine  
guanidine derivative; aminoguanidine; guanidine hydrochloride; desmin;  
glycerol; pyrene; oligomer; unclassified drug

**MEDICAL DESCRIPTORS:**

\*protein structure  
structure analysis; protein targeting; protein binding; point mutation;  
circular dichroism; fluorescence; sedimentation; protein secondary  
structure; solubilization; protein tertiary structure; protein quaternary  
structure; nonhuman; article; priority journal

DRUG TERMS (UNCONTROLLED): alpha b crystallin

CAS REGISTRY NO.: 1119-34-2, 15595-35-4, 7004-12-8, 74-79-3 (arginine);  
1068-42-4, 2582-30-1, 79-17-4 (aminoguanidine); 50-01-1 (guanidine)

hydrochloride); 56-81-5 (glycerol); 129-00-0 (pyrene)  
SECTION HEADINGS:

029 Clinical and Experimental Biochemistry

**Structural perturbation and enhancement of the chaperone-like activity of alpha-crystallin by arginine hydrochloride**

Structural perturbation of alpha-crystallin is shown to enhance its molecular chaperone-like activity in preventing aggregation of target proteins...

...and negatively charged side-chains, increases the chaperone-like activity of calf eye lens alpha-crystallin as well as recombinant human alphaA- and alphaB-crystallins. Arginine-induced increase in the chaperone activity is more pronounced for alphaB-crystallin than for alphaA-crystallin. Other guanidinium compounds such as aminoguanidine hydrochloride and guanidine hydrochloride also show a similar effect, but to different extents. A point mutation, R120G, in alphaB-crystallin that is associated with desmin-related myopathy, results in a significant loss of chaperone-like...

...considerable extent. We have investigated the effect of arginine on the structural changes of alpha-crystallin by circular dichroism, fluorescence, and glycerol gradient sedimentation. Far-UV CD spectra show no significant...

...presence of arginine. Glycerol gradient sedimentation shows a significant decrease in the size of alpha-crystallin oligomer in the presence of arginine. Increased exposure of hydrophobic surfaces of alpha-crystallin, as monitored by pyrene-solubilization and ANS-fluorescence, is observed in the presence of arginine...

...subtle changes in the tertiary structure and significant changes in the quaternary structure of alpha-crystallin and enhances its chaperone-like activity significantly. This study should prove useful in designing strategies...

DRUG DESCRIPTORS:

\*chaperone; \*alpha crystallin --endogenous compound--ec; \*arginine

DRUG TERMS (UNCONTROLLED): alpha b crystallin

7/9,K/9 (Item 1 from file: 155)

DIALOG(R) File 155:MEDLINE(R)

(c) format only 2006 Dialog. All.rts. reserv.

20362892 PMID: 15827547

Arginine hydrochloride enhances the dynamics of subunit assembly and the chaperone-like activity of alpha-crystallin.

Srinivas V; Raman B; Rao K Sridhar; Ramakrishna T; Rao Ch Mohan  
Centre for Cellular and Molecular Biology, Hyderabad, India.

Molecular vision electronic resource (United States) 2005, 11  
p249-55, ISSN 1090-0535--Electronic Journal Code: 9605351

Publishing Model Electronic

Document type: Journal Article

Languages: ENGLISH

Main Citation Owner: NLM

Record type: MEDLINE; Completed

Subfile: INDEX MEDICUS

**PURPOSE:** Alpha- **crystallin** , a major eye lens protein, bears homology with small heat shock proteins (sHsps) and exhibits molecular chaperone-like activity. Structural perturbation by temperature or low concentrations of denaturants leads to enhancement of its chaperone-like activity. We have earlier demonstrated similar enhancement of chaperone-like activity using biologically compatible solutes such as **arginine hydrochloride** and aminoguanidine. The purpose of the present study is to get an insight into the mechanism of the arginine induced enhancement of chaperone-like activity of alpha- **crystallin** . **METHODS:** The effect of **arginine hydrochloride** on the chaperone-like activity of alpha- **crystallin** at 25 degrees C was studied using DTT induced aggregation of insulin as a model system. Changes in the accessibility of the thiol group near the end of the alpha- **crystallin** domain in the absence and the presence of **arginine hydrochloride** were studied using dithiobisnitrobenzoic acid. Fluorescence resonance energy transfer studies were performed to investigate changes in the dynamics of the subunit assembly. Urea induced denaturation studies of alpha- **crystallin** were carried out to investigate structural destabilization of alpha- **crystallin** , if any, in the presence of **arginine hydrochloride** . **RESULTS:** **Arginine hydrochloride** increases the chaperone-like activity of alpha- **crystallin** several fold towards DTT induced aggregation of insulin at room temperature. Our study shows that both the extent and the rate of accessibility of the thiol group are increased in the presence of arginine. Fluorescence resonance energy transfer experiments show that **arginine hydrochloride** significantly increases the subunit exchange between the oligomers of alpha- **crystallin** . Arginine induced structural perturbation and loosening of subunit assembly of alpha- **crystallin** leads to overall destabilization of the protein as reflected by the urea denaturation study. **CONCLUSIONS:** Arginine perturbs the tertiary and quaternary structure of alpha- **crystallin** and enhances the dynamics of the subunit assembly leading to enhanced chaperone-like activity. Thus, in addition to size, surface hydrophobicity, and charge distribution, the dynamics of the subunit assembly appears to be one of the critical factors that can modulate the chaperone activity.

**Descriptors:** \*Arginine--pharmacology--PD; \*Molecular Chaperones --metabolism--ME; \*alpha-Crystallins--drug effects--DE; Animals; Cattle; Disulfides; Dithiothreitol; Fluorescent Dyes; Lens, Crystalline--chemistry --CH; Protein Subunits--chemistry--CH; Protein Subunits--metabolism--ME; Recombinant Proteins--chemistry--CH; Recombinant Proteins--drug effects --DE; Recombinant Proteins--metabolism--ME; Solubility; Spectrometry, Fluorescence; alpha-Crystallins--chemistry--CH; alpha-Crystallins --metabolism--ME

CAS Registry No.: 0 (Disulfides); 0 (Fluorescent Dyes); 0 (Molecular Chaperones); 0 (Protein Subunits); 0 (Recombinant Proteins); 0 (alpha-Crystallins); 3483-12-3 (Dithiothreitol); 74-79-3 (Arginine)

Record Date Created: 20050413

Record Date Completed: 20060413

Date of Electronic Publication: 20050401

**Arginine hydrochloride enhances the dynamics of subunit assembly and the chaperone-like activity of alpha- crystallin .**

**PURPOSE:** Alpha- **crystallin** , a major eye lens protein, bears homology with small heat shock proteins (sHsps) and exhibits...

...have earlier demonstrated similar enhancement of chaperone-like activity using biologically compatible solutes such as **arginine hydrochloride** and aminoguanidine. The purpose of the present study is to get an insight

into the mechanism of the arginine induced enhancement of chaperone-like activity of alpha- **crystallin** . METHODS: The effect of **arginine hydrochloride** on the chaperone-like activity of alpha- **crystallin** at 25 degrees C was studied using DTT induced aggregation of insulin as a model system. Changes in the accessibility of the thiol group near the end of the alpha- **crystallin** domain in the absence and the presence of **arginine hydrochloride** were studied using dithiobisnitrobenzoic acid. Fluorescence resonance energy transfer studies were performed to investigate changes in the dynamics of the subunit assembly. Urea induced denaturation studies of alpha- **crystallin** were carried out to investigate structural destabilization of alpha- **crystallin** , if any, in the presence of **arginine hydrochloride** . RESULTS: **Arginine hydrochloride** increases the chaperone-like activity of alpha- **crystallin** several fold towards DTT induced aggregation of insulin at room temperature. Our study shows that...

... group are increased in the presence of arginine. Fluorescence resonance energy transfer experiments show that **arginine hydrochloride** significantly increases the subunit exchange between the oligomers of alpha- **crystallin** . Arginine induced structural perturbation and loosening of subunit assembly of alpha- **crystallin** leads to overall destabilization of the protein as reflected by the urea denaturation study. CONCLUSIONS: Arginine perturbs the tertiary and quaternary structure of alpha- **crystallin** and enhances the dynamics of the subunit assembly leading to enhanced chaperone-like activity. Thus...

7/9,K/10 (Item 2 from file: 155)

DIALOG(R) File 155:MEDLINE(R)

(c) format only 2006 Dialog. All rts. reserv.

14316050 PMID: 12761397

**Structural perturbation and enhancement of the chaperone-like activity of alpha- crystallin by arginine hydrochloride .**

Srinivas Volety; Raman Bakthisaran; Rao Kunchala Sridhar; Ramakrishna Tangirala; Rao Ch Mohan

Centre for Cellular & Molecular Biology, Hyderabad 500 007, India.

Protein science - a publication of the Protein Society (United States)

Jun 2003, 12 (6) p1262-70, ISSN 0961-8368--Print Journal Code: 9211750

Publishing Model Print

Document type: Journal Article

Languages: ENGLISH

Main Citation Owner: NLM

Record type: MEDLINE; Completed

Subfile: INDEX MEDICUS

Structural perturbation of alpha- **crystallin** is shown to enhance its molecular chaperone-like activity in preventing aggregation of target proteins. We demonstrate that arginine, a biologically compatible molecule that is known to bind to the peptide backbone and negatively charged side-chains, increases the chaperone-like activity of calf eye lens alpha- **crystallin** as well as recombinant human alphaA- and alphaB-crystallins. Arginine-induced increase in the chaperone activity is more pronounced for alphaB- **crystallin** than for alphaA- **crystallin** . Other guanidinium compounds such as aminoguanidine hydrochloride and guanidine hydrochloride also show a similar effect, but to different extents. A point mutation, R120G, in alphaB- **crystallin** that is associated with desmin-related myopathy, results in a significant loss of chaperone-like activity.

Arginine restores the activity of mutant protein to a considerable extent. We have investigated the effect of arginine on the structural changes of alpha- **crystallin** by circular dichroism, fluorescence, and glycerol gradient sedimentation. Far-UV CD spectra show no significant changes in secondary structure, whereas near-UV CD spectra show subtle changes in the presence of arginine. Glycerol gradient sedimentation shows a significant decrease in the size of alpha- **crystallin** oligomer in the presence of arginine. Increased exposure of hydrophobic surfaces of alpha- **crystallin** , as monitored by pyrene-solubilization and ANS-fluorescence, is observed in the presence of arginine. These results show that arginine brings about subtle changes in the tertiary structure and significant changes in the quaternary structure of alpha- **crystallin** and enhances its chaperone-like activity significantly. This study should prove useful in designing strategies to improve chaperone function for therapeutic applications.

Descriptors: \*Arginine--pharmacology--PD; \*Crystallins--chemistry--CH; Animals; Cattle; Centrifugation, Density Gradient; Circular Dichroism; Crystallins--metabolism--ME; Dithiothreitol; Guanidine--pharmacology--PD; Insulin--chemistry--CH; Insulin--metabolism--ME; Protein Conformation --drug effects--DE; Pyrenes--chemistry--CH; Solubility; Spectrometry, Fluorescence; Time Factors

CAS Registry No.: 0 (Crystallins); 0 (Pyrenes); 11061-68-0 (Insulin); 113-00-8 (Guanidine); 129-00-0 (pyrene); 3483-12-3 (Dithiothreitol); 74-79-3 (Arginine)

Record Date Created: 20030522

Record Date Completed: 20041005

**Structural perturbation and enhancement of the chaperone-like activity of alpha- crystallin by arginine hydrochloride .**

Structural perturbation of alpha- **crystallin** is shown to enhance its molecular chaperone-like activity in preventing aggregation of target proteins...

... and negatively charged side-chains, increases the chaperone-like activity of calf eye lens alpha- **crystallin** as well as recombinant human alphaA- and alphaB-crystallins. Arginine-induced increase in the chaperone activity is more pronounced for alphaB- **crystallin** than for alphaA- **crystallin** . Other guanidinium compounds such as aminoguanidine hydrochloride and guanidine hydrochloride also show a similar effect, but to different extents. A point mutation, R120G, in alphaB- **crystallin** that is associated with desmin-related myopathy, results in a significant loss of chaperone-like...

... considerable extent. We have investigated the effect of arginine on the structural changes of alpha- **crystallin** by circular dichroism, fluorescence, and glycerol gradient sedimentation. Far-UV CD spectra show no significant...

... presence of arginine. Glycerol gradient sedimentation shows a significant decrease in the size of alpha- **crystallin** oligomer in the presence of arginine. Increased exposure of hydrophobic surfaces of alpha- **crystallin** , as monitored by pyrene-solubilization and ANS-fluorescence, is observed in the presence of arginine...

... subtle changes in the tertiary structure and significant changes in the quaternary structure of alpha- **crystallin** and enhances its chaperone-like activity significantly. This study should prove useful in designing strategies...

? s arginine and hydrochloride  
416125 ARGININE  
382785 HYDROCHLORIDE  
S8 4708 ARGININE AND HYDROCHLORIDE  
? s s1 and s8  
40380 S1  
4708 S8  
S9 20 S1 AND S8  
? t s9/9,k/1-5

9/9,K/1 (Item 1 from file: 5)  
DIALOG(R) File 5:Biosis Previews(R)  
(c) 2007 The Thomson Corporation. All rts. reserv.

0016021600 BIOSIS NO.: 200600366995

**Effect of site-directed mutagenesis of methylglyoxal-modifiable arginine residues on the structure and chaperone function of human alpha A-crystallin**

AUTHOR: Biswas Ashis; Miller Antonia; Oya-Ito Tomoko; Santhoshkumar Puttur; Bhat Manjunatha; Nagaraj Ram H (Reprint)

AUTHOR ADDRESS: Case Western Reserve Univ, Dept Ophthalmol, Cleveland, OH 44106 USA\*\*USA

AUTHOR E-MAIL ADDRESS: ram.nagaraj@case.edu

JOURNAL: Biochemistry 45 (14): p4569-4577 APR 11 2006 2006

ISSN: 0006-2960

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

**ABSTRACT:** We reported previously that chemical modification of human alpha A- **crystallin** by a metabolic dicarbonyl compound, methylglyoxal (MGO), enhances its chaperone-like function, a phenomenon which we attributed to formation of argpyrimidine at **arginine** residues (R) 21 49, and 103. This structural change removes the positive charge on the **arginine** residues. To explore this mechanism further, we replaced these three R residues with a neutral alanine (A) residue one at a time or in combination and examined the impact on the structure and chaperone function. Measurement of intrinsic tryptophan fluorescence and near-UV CD spectra revealed alteration of the microenvironment of aromatic amino acid residues in mutant proteins. When compared to wild-type (wt) alpha A- **crystallin**, the chaperone function of R21A and R103A mutants increased 20% and 18% as measured by the insulin aggregation assay and increased it as much as 39% and 28% when measured by the citrate synthase (CS) aggregation assay. While the R49A mutant lost most of its chaperone function, R21A/R103A and R21A/R49A/R103A mutants had slightly better function (6-14% and 10-14%) than the wt protein in these assays. R21A and R103A mutants had higher surface hydrophobicity than wt alpha A- **crystallin**, but the R49A mutant had lower hydrophobicity. R21A and R103A mutants, but not the R49A mutant, were more efficient than wt protein in refolding guanidine **hydrochloride** -treated malate dehydrogenase to its native state. Our findings indicate that the positive charges on R21, R49, and R103 are important determinants of the chaperone function of alpha A- **crystallin** and suggest that chemical modification of **arginine** residues may play a role in protein aggregation during lens aging and cataract formation.

REGISTRY NUMBERS: 78-98-8: methylglyoxal

DESCRIPTORS:

MAJOR CONCEPTS: Biochemistry and Molecular Biophysics

BIOSYSTEMATIC NAMES: Enterobacteriaceae--Facultatively Anaerobic

Gram-Negative Rods, Eubacteria, Bacteria, Microorganisms

ORGANISMS: Escherichia coli (Enterobacteriaceae)--expression system

COMMON TAXONOMIC TERMS: Bacteria; Eubacteria; Microorganisms

CHEMICALS & BIOCHEMICALS: methylglyoxal {MGO}; alpha-A- **crystallin** --  
methylglyoxal-modifiable **arginine** residue

METHODS & EQUIPMENT: site-directed mutagenesis--laboratory techniques,  
genetic techniques

CONCEPT CODES:

10060 Biochemistry studies - General

31000 Physiology and biochemistry of bacteria

BIOSYSTEMATIC CODES:

06702 Enterobacteriaceae

**Effect of site-directed mutagenesis of methylglyoxal-modifiable arginine residues on the structure and chaperone function of human alpha A-crystallin**

ABSTRACT: We reported previously that chemical modification of human alpha A- **crystallin** by a metabolic dicarbonyl compound, methylglyoxal (MGO), enhances its chaperone-like function, a phenomenon which we attributed to formation of argpyrimidine at **arginine** residues (R) 21 49, and 103. This structural change removes the positive charge on the **arginine** residues. To explore this mechanism further, we replaced these three R residues with a neutral...

...aromatic amino acid residues in mutant proteins. When compared to wild-type (wt) alpha A- **crystallin**, the chaperone function of R21A and R103A mutants increased 20% and 18% as measured by...

...in these assays. R21A and R103A mutants had higher surface hydrophobicity than wt alpha A- **crystallin**, but the R49A mutant had lower hydrophobicity. R21A and R103A mutants, but not the R49A mutant, were more efficient than wt protein in refolding guanidine **hydrochloride**-treated malate dehydrogenase to its native state. Our findings indicate that the positive charges on R21, R49, and R103 are important determinants of the chaperone function of alpha A- **crystallin** and suggest that chemical modification of **arginine** residues may play a role in protein aggregation during lens aging and cataract formation.

DESCRIPTORS:

CHEMICALS & BIOCHEMICALS: ...alpha-A- **crystallin** --....

...methylglyoxal-modifiable **arginine** residue

9/9,K/2 (Item 2 from file: 5)

DIALOG(R) File 5:Biosis Previews(R)

(c) 2007 The Thomson Corporation. All rts. reserv.

0015321550 BIOSIS NO.: 200510016050

**Arginine hydrochloride enhances the dynamics of subunit assembly and the chaperone-like activity of alpha- crystallin**

AUTHOR: Srinivas V; Raman B; Rao K Sridhar; Ramakrishna T; Rao Ch Mohan  
(Reprint)

AUTHOR ADDRESS: Ctr Cellular and Mol Biol, Uppal Rd, Hyderabad 500007,  
Andhra Pradesh, India\*\*India

AUTHOR E-MAIL ADDRESS: mohan@ccmb.res.in  
JOURNAL: Molecular Vision 11 (27-29): p249-255 APR 1 05 2005  
ISSN: 1090-0535  
DOCUMENT TYPE: Article  
RECORD TYPE: Abstract  
LANGUAGE: English

**ABSTRACT:** Purpose: alpha- **Crystallin** , a major eye lens protein, bears homology with small heat shock proteins (sHsps) and exhibits molecular chaperone-like activity. Structural perturbation by temperature or low concentrations of denaturants leads to enhancement of its chaperone-like activity. We have earlier demonstrated similar enhancement of chaperone-like activity using biologically compatible solutes such as **arginine hydrochloride** and aminoguanidine. The purpose of the present study is to get an insight into the mechanism of the **arginine** induced enhancement of chaperone-like activity of alpha- **crystallin** .Methods: The effect of **arginine hydrochloride** on the chaperone-like activity of alpha- **crystallin** at 25 degrees C was studied using DTT induced aggregation of insulin as a model system. Changes in the accessibility of the thiol group near the end of the alpha- **crystallin** domain in the absence and the presence of **arginine hydrochloride** were studied using dithiobisnitrobenzoic acid. Fluorescence resonance energy transfer studies were performed to investigate changes in the dynamics of the subunit assembly. Urea induced denaturation studies of alpha- **crystallin** were carried out to investigate structural destabilization of alpha- **crystallin** , if any, in the presence of **arginine hydrochloride** .Results: **Arginine hydrochloride** increases the chaperone-like activity of alpha- **crystallin** several fold towards DTT induced aggregation of insulin at room temperature. Our study shows that both the extent and the rate of accessibility of the thiol group are increased in the presence of **arginine** . Fluorescence resonance energy transfer experiments show that **arginine hydrochloride** significantly increases the subunit exchange between the oligomers of alpha- **crystallin** . **Arginine** induced structural perturbation and loosening of subunit assembly of alpha- **crystallin** leads to overall destabilization of the protein as reflected by the urea denaturation study.Conclusions: **Arginine** perturbs the tertiary and quaternary structure of alpha- **crystallin** and enhances the dynamics of the subunit assembly leading to enhanced chaperone-like activity. Thus, in addition to size, surface hydrophobicity, and charge distribution, the dynamics of the subunit assembly appears to be one of the critical factors that can modulate the chaperone activity.

REGISTRY NUMBERS: 9004-10-8: insulin; 79-17-4: aminoguanidine; 32042-43-6: **arginine hydrochloride**

**DESCRIPTORS:**

MAJOR CONCEPTS: Biochemistry and Molecular Biophysics; Sense Organs--  
Sensory Reception

ORGANISMS: PARTS ETC: eye lens

CHEMICALS & BIOCHEMICALS: insulin; small heat shock proteins;  
aminoguanidine; **arginine hydrochloride** ; alpha- **crystallin** --  
chaperone-like activity

**CONCEPT CODES:**

10060 Biochemistry studies - General

10064 Biochemistry studies - Proteins, peptides and amino acids

20004 Sense organs - Physiology and biochemistry



**Arginine hydrochloride enhances the dynamics of subunit assembly and the chaperone-like activity of alpha- crystallin**

**ABSTRACT:** Purpose: alpha- **Crystallin** , a major eye lens protein, bears homology with small heat shock proteins (sHsps) and exhibits...

...have earlier demonstrated similar enhancement of chaperone-like activity using biologically compatible solutes such as **arginine hydrochloride** and aminoguanidine. The purpose of the present study is to get an insight into the mechanism of the **arginine** induced enhancement of chaperone-like activity of alpha- **crystallin** .Methods: The effect of **arginine hydrochloride** on the chaperone-like activity of alpha- **crystallin** at 25 degrees C was studied using DTT induced aggregation of insulin as a model system. Changes in the accessibility of the thiol group near the end of the alpha- **crystallin** domain in the absence and the presence of **arginine hydrochloride** were studied using dithiobisnitrobenzoic acid. Fluorescence resonance energy transfer studies were performed to investigate changes in the dynamics of the subunit assembly. Urea induced denaturation studies of alpha- **crystallin** were carried out to investigate structural destabilization of alpha- **crystallin** , if any, in the presence of **arginine hydrochloride** .Results: **Arginine hydrochloride** increases the chaperone-like activity of alpha- **crystallin** several fold towards DTT induced aggregation of insulin at room temperature. Our study shows that...

...and the rate of accessibility of the thiol group are increased in the presence of **arginine** . Fluorescence resonance energy transfer experiments show that **arginine hydrochloride** significantly increases the subunit exchange between the oligomers of alpha- **crystallin** . **Arginine** induced structural perturbation and loosening of subunit assembly of alpha- **crystallin** leads to overall destabilization of the protein as reflected by the urea denaturation study.Conclusions: **Arginine** perturbs the tertiary and quaternary structure of alpha- **crystallin** and enhances the dynamics of the subunit assembly leading to enhanced chaperone-like activity. Thus...

...REGISTRY NUMBERS: **arginine hydrochloride**

DESCRIPTORS:

CHEMICALS & BIOCHEMICALS: ... **arginine hydrochloride** ; ...

...alpha- **crystallin** --

9/9,K/3 (Item 3 from file: 5)

DIALOG(R)File 5:Biosis Previews(R)

(c) 2007 The Thomson Corporation. All rts. reserv.

0014361712 BIOSIS NO.: 200300320431

**Structural perturbation and enhancement of the chaperone-like activity of alpha- crystallin by arginine hydrochloride .**

AUTHOR: Srinivas Volety; Raman Bakthisaran; Rao Kunchala Sridhar; Ramakrishna Tangirala; Rao Ch Mohan (Reprint)

AUTHOR ADDRESS: Centre for Cellular and Molecular Biology, Uppal Road, Hyderabad, 500 007, India\*\*India

AUTHOR E-MAIL ADDRESS: mohan@ccmb.res.in

JOURNAL: Protein Science 12 (6): p1262-1270 June 2003 2003

MEDIUM: print

ISSN: 0961-8368

DOCUMENT TYPE: Article  
RECORD TYPE: Abstract  
LANGUAGE: English

ABSTRACT: Structural perturbation of alpha- **crystallin** is shown to enhance its molecular chaperone-like activity in preventing aggregation of target proteins. We demonstrate that **arginine**, a biologically compatible molecule that is known to bind to the peptide backbone and negatively charged side-chains, increases the chaperone-like activity of calf eye lens alpha- **crystallin** as well as recombinant human alphaA- and alphaB-crystallins. **Arginine**-induced increase in the chaperone activity is more pronounced for alphaB- **crystallin** than for alphaA- **crystallin**. Other guanidinium compounds such as aminoguanidine **hydrochloride** and guanidine **hydrochloride** also show a similar effect, but to different extents. A point mutation, R120G, in alphaB- **crystallin** that is associated with desmin-related myopathy, results in a significant loss of chaperone-like activity. **Arginine** restores the activity of mutant protein to a considerable extent. We have investigated the effect of **arginine** on the structural changes of alpha- **crystallin** by circular dichroism, fluorescence, and glycerol gradient sedimentation. Far-UV CD spectra show no significant changes in secondary structure, whereas near-UV CD spectra show subtle changes in the presence of **arginine**. Glycerol gradient sedimentation shows a significant decrease in the size of alpha- **crystallin** oligomer in the presence of **arginine**. Increased exposure of hydrophobic surfaces of alpha- **crystallin**, as monitored by pyrene-solubilization and ANS-fluorescence, is observed in the presence of **arginine**. These results show that **arginine** brings about subtle changes in the tertiary structure and significant changes in the quaternary structure of alpha- **crystallin** and enhances its chaperone-like activity significantly. This study should prove useful in designing strategies to improve chaperone function for therapeutic applications.

REGISTRY NUMBERS: 1119-34-2Q: **arginine hydrochloride**; 15595-35-4Q: **arginine hydrochloride**; 32042-43-6Q: **arginine hydrochloride**; 1937-19-5Q: aminoguanidine **hydrochloride**; 16139-18-7Q: aminoguanidine **hydrochloride**

DESCRIPTORS:

MAJOR CONCEPTS: Biochemistry and Molecular Biophysics

BIOSYSTEMATIC NAMES: Bovidae--Artiodactyla, Mammalia, Vertebrata, Chordata, Animalia; Hominidae--Primates, Mammalia, Vertebrata, Chordata, Animalia

ORGANISMS: cow (Bovidae); human (Hominidae)

ORGANISMS: PARTS ETC: lens--sensory system

COMMON TAXONOMIC TERMS: Artiodactyls; Nonhuman Vertebrates; Nonhuman Mammals; Animals; Chordates; Humans; Mammals; Primates; Vertebrates

DISEASES: desmin-related myopathy--muscle disease

CHEMICALS & BIOCHEMICALS: alpha- **crystallin** --structure, chaperone-like activity; **arginine hydrochloride**; alpha-A **crystallin**; alpha-B **crystallin**; aminoguanidine **hydrochloride**

METHODS & EQUIPMENT: far-UV circular dichroism spectroscopy--laboratory techniques, spectrum analysis techniques; near-UV circular dichroism spectroscopy--laboratory techniques, spectrum analysis techniques; fluorescence assay--laboratory techniques, spectrum analysis techniques; glycerol gradient sedimentation--laboratory techniques

MISCELLANEOUS TERMS: drug development

CONCEPT CODES:

10060 Biochemistry studies - General  
17506 Muscle - Pathology  
20004 Sense organs - Physiology and biochemistry  
BIOSYSTEMATIC CODES:  
85715 Bovidae  
86215 Hominidae

**Structural perturbation and enhancement of the chaperone-like activity of  
alpha- crystallin by arginine hydrochloride .**

**ABSTRACT:** Structural perturbation of alpha- **crystallin** is shown to enhance its molecular chaperone-like activity in preventing aggregation of target proteins. We demonstrate that **arginine** , a biologically compatible molecule that is known to bind to the peptide backbone and negatively charged side-chains, increases the chaperone-like activity of calf eye lens alpha- **crystallin** as well as recombinant human alphaA- and alphaB-crystallins. **Arginine** -induced increase in the chaperone activity is more pronounced for alphaB- **crystallin** than for alphaA- **crystallin** . Other guanidinium compounds such as aminoguanidine **hydrochloride** and guanidine **hydrochloride** also show a similar effect, but to different extents. A point mutation, R120G, in alphaB- **crystallin** that is associated with desmin-related myopathy, results in a significant loss of chaperone-like activity. **Arginine** restores the activity of mutant protein to a considerable extent. We have investigated the effect of **arginine** on the structural changes of alpha- **crystallin** by circular dichroism, fluorescence, and glycerol gradient sedimentation. Far-UV CD spectra show no significant...

...in secondary structure, whereas near-UV CD spectra show subtle changes in the presence of **arginine** . Glycerol gradient sedimentation shows a significant decrease in the size of alpha- **crystallin** oligomer in the presence of **arginine** . Increased exposure of hydrophobic surfaces of alpha- **crystallin** , as monitored by pyrene-solubilization and ANS-fluorescence, is observed in the presence of **arginine** . These results show that **arginine** brings about subtle changes in the tertiary structure and significant changes in the quaternary structure of alpha- **crystallin** and enhances its chaperone-like activity significantly. This study should prove useful in designing strategies...

...REGISTRY NUMBERS: **arginine hydrochloride** ; ...

... **arginine hydrochloride** ; ...

... **arginine hydrochloride** ; ...

...aminoguanidine **hydrochloride** ; ...

...aminoguanidine **hydrochloride**

**DESCRIPTORS:**

**CHEMICALS & BIOCHEMICALS:** alpha- **crystallin** --....

... **arginine hydrochloride** ; ...

...alpha-A **crystallin** ; ...

...alpha-B **crystallin** ; ...

...aminoguanidine **hydrochloride**

9/9,K/4 (Item 1 from file: 34)

DIALOG(R) File 34:SciSearch(R) Cited Ref Sci

(c) 2007 The Thomson Corp. All rts. reserv.

15023860 Genuine Article#: 031GH Number of References: 59

Title: **Effect of site-directed mutagenesis of methylglyoxal-modifiable arginine residues on the structure and chaperone function of human alpha A- crystallin**

Author(s): Biswas A; Miller A; Oya-Ito T; Santhoshkumar P; Bhat M; Nagaraj RH (REPRINT)

Corporate Source: Case Western Reserve Univ, Dept Ophthalmol, Cleveland//OH/44106 (REPRINT); Case Western Reserve Univ, Dept Ophthalmol, Cleveland//OH/44106; Case Western Reserve Univ, Dept Pharmacol, Cleveland//OH/44106; Cleveland Clin Fdn, Ctr Anesthesiol Res, Cleveland//OH/44195; Univ Missouri, Mason Eye Inst, Columbia//MO/65212 (ram.nagaraj@case.edu)

Journal: BIOCHEMISTRY, 2006, V45, N14 (APR 11), P4569-4577

ISSN: 0006-2960 Publication date: 20060411

Publisher: AMER CHEMICAL SOC, 1155 16TH ST, NW, WASHINGTON, DC 20036 USA

Language: English Document Type: ARTICLE

Geographic Location: USA

Journal Subject Category: BIOCHEMISTRY & MOLECULAR BIOLOGY

Abstract: We reported previously that chemical modification of human alpha A- **crystallin** by a metabolic dicarbonyl compound, methylglyoxal (MGO), enhances its chaperone-like function, a phenomenon which we attributed to formation of argpyrimidine at **arginine** residues (R) 21 49, and 103. This structural change removes the positive charge on the **arginine** residues. To explore this mechanism further, we replaced these three R residues with a neutral alanine (A) residue one at a time or in combination and examined the impact on the structure and chaperone function. Measurement of intrinsic tryptophan fluorescence and near-UV CD spectra revealed alteration of the microenvironment of aromatic amino acid residues in mutant proteins. When compared to wild-type (wt) alpha A- **crystallin**, the chaperone function of R21A and R103A mutants increased 20% and 18% as measured by the insulin aggregation assay and increased it as much as 39% and 28% when measured by the citrate synthase (CS) aggregation assay. While the R49A mutant lost most of its chaperone function, R21A/R103A and R21A/R49A/R103A mutants had slightly better function (6-14% and 10-14%) than the wt protein in these assays. R21A and R103A mutants had higher surface hydrophobicity than wt alpha A- **crystallin**, but the R49A mutant had lower hydrophobicity. R21A and R103A mutants, but not the R49A mutant, were more efficient than wt protein in refolding guanidine **hydrochloride**-treated malate dehydrogenase to its native state. Our findings indicate that the positive charges on R21, R49, and R103 are important determinants of the chaperone function of alpha A- **crystallin** and suggest that chemical modification of **arginine** residues may play a role in protein aggregation during lens aging and cataract formation.

Identifiers--Keyword Plus(R): HEAT-SHOCK-PROTEIN; HUMAN LENS PROTEINS; B-CRYSTALLIN; MOLECULAR CHAPERONE; MAILLARD REACTION; CROSS-LINKS; POSTTRANSLATIONAL MODIFICATIONS; OLIGOMERIC SIZE; SERUM-ALBUMIN; GLYCATION

Cited References:

AHMED MU, 1997, V324, P565, BIOCHEM J 2

AHMED N, 2003, V44, P5287, INVEST OPHTH VIS SCI

BAYNES JW, 2001, V36, P1527, EXP GERONTOL  
 BERA S, 2002, V41, P12421, BIOCHEMISTRY-US  
 BHATTACHARYYA J, 1998, V46, P249, BIOCHEM MOL BIOL INT  
 BHATTACHARYYA J, 2002, V21, P65, J PROTEIN CHEM  
 BIEMEL KM, 2002, V277, P24907, J BIOL CHEM  
 BISWAS A, 2004, V279, P42648, J BIOL CHEM  
 BLOEMENDAL H, 1982, V12, P1, CRC CRIT REV BIOCH  
 BLOEMENDAL H, 2004, V86, P407, PROG BIOPHYS MOL BIO  
 BOVA MP, 1999, V96, P6137, P NATL ACAD SCI USA  
 CHELLAN P, 1999, V368, P98, ARCH BIOCHEM BIOPHYS  
 CHERIAN M, 1995, V208, P675, BIOCHEM BIOPH RES CO  
 CLARK JI, 2004, V45, INVEST OPHTHALMOL VI  
 DAS KP, 1995, V369, P321, FEBS LETT  
 DEJONG WW, 1998, V22, P151, INT J BIOL MACROMOL  
 DEJONG WW, 1993, V10, P103, MOL BIOL EVOL  
 DERHAM BK, 1999, V18, P463, PROG RETIN EYE RES  
 FUJII N, 2001, V74, P477, PHOTOCHEM PHOTOBIO  
 GUPTA R, 2004, V279, P44258, J BIOL CHEM  
 HAIK GM, 1994, V59, P497, EXP EYE RES  
 HORWITZ J, 1992, V89, P10449, P NATL ACAD SCI USA  
 HORWITZ J, 2003, V76, P145, EXP EYE RES  
 ITO H, 2001, V276, P5346, J BIOL CHEM  
 IWAKI T, 1990, V38, P31, J HISTOCHEM CYTOCHEM  
 KAPPAHN RJ, 2003, V42, P15310, BIOCHEMISTRY-US  
 KUMAR MS, 2004, V379, P273, BIOCHEM J 2  
 KUMAR MS, 2005, V280, P21726, J BIOL CHEM  
 KUMAR LVS, 1999, V274, P24137, J BIOL CHEM  
 LIAO JH, 2002, V295, P854, BIOCHEM BIOPH RES CO  
 LO TWC, 1994, V269, P32299, J BIOL CHEM  
 LYONS TJ, 1991, V40, P1010, DIABETES  
 MCCLURE WO, 1966, V5, P1908, BIOCHEMISTRY-US  
 MUCHOWSKI PJ, 1998, V95, P1004, P NATL ACAD SCI USA  
 NAGARAJ RH, 1996, V382, P234, FEBS LETT  
 NAGARAJ RH, 1991, V88, P10257, P NATL ACAD SCI USA  
 NAGARAJ RH, 2003, V42, P10746, BIOCHEMISTRY-US  
 PADAYATTI PS, 2001, V42, P1299, INVEST OPHTH VIS SCI  
 PASTA SY, 2003, V278, P51159, J BIOL CHEM  
 PASTA SY, 2004, V10, P655, MOL VIS  
 PROVENCHER SW, 1981, V20, P33, BIOCHEMISTRY-US  
 RAMAN B, 1995, V365, P133, FEBS LETT  
 RAWAT U, 1998, V273, P9415, J BIOL CHEM  
 REDDY GB, 2000, V275, P4565, J BIOL CHEM  
 SAHA S, 2004, V57, P610, PROTEINS  
 SANTHOSHKUMAR P, 2001, V276, P47094, J BIOL CHEM  
 SHIPANOVA IN, 1997, V344, P29, ARCH BIOCHEM BIOPHYS  
 SHROFF NP, 2001, V220, P127, MOL CELL BIOCHEM  
 SPECTOR A, 1971, V124, P337, BIOCHEM J  
 SREELAKSHMI Y, 2005, V44, P12245, BIOCHEMISTRY-US  
 SUN TX, 1997, V272, P6220, J BIOL CHEM  
 THORNALLEY PJ, 1993, V14, P287, MOL ASPECTS MED  
 THORNALLEY PJ, 2003, V375, P581, BIOCHEM J 3  
 VANBOEKEL MAM, 1996, V28, P32, OPHTHALMIC RES S1  
 WANG KY, 2000, V267, P4705, EUR J BIOCHEM  
 WANG KY, 1994, V269, P13601, J BIOL CHEM  
 WELLSKNECHT KJ, 1996, V11, P41, NEPHROL DIAL TRAN S5  
 WILKER SC, 2001, V290, P353, ANAL BIOCHEM  
 WYATT PJ, 1993, V272, P1, ANAL CHIM ACTA

**Title: Effect of site-directed mutagenesis of methylglyoxal-modifiable arginine residues on the structure and chaperone function of human alpha A- crystallin**

**Abstract:** We reported previously that chemical modification of human alpha A- **crystallin** by a metabolic dicarbonyl compound, methylglyoxal (MGO), enhances its chaperone-like function, a phenomenon which we attributed to formation of argpyrimidine at **arginine** residues (R) 21 49, and 103. This structural change removes the positive charge on the **arginine** residues. To explore this mechanism further, we replaced these three R residues with a neutral...

...aromatic amino acid residues in mutant proteins. When compared to wild-type (wt) alpha A- **crystallin**, the chaperone function of R21A and R103A mutants increased 20% and 18% as measured by...

...in these assays. R21A and R103A mutants had higher surface hydrophobicity than wt alpha A- **crystallin**, but the R49A mutant had lower hydrophobicity. R21A and R103A mutants, but not the R49A mutant, were more efficient than wt protein in refolding guanidine **hydrochloride**-treated malate dehydrogenase to its native state. Our findings indicate that the positive charges on R21, R49, and R103 are important determinants of the chaperone function of alpha A- **crystallin** and suggest that chemical modification of **arginine** residues may play a role in protein aggregation during lens aging and cataract formation.

...Identifiers--HEAT-SHOCK-PROTEIN; HUMAN LENS PROTEINS; B- **CRYSTALLIN**; MOLECULAR CHAPERONE; MAILLARD REACTION; CROSS-LINKS; POSTTRANSLATIONAL MODIFICATIONS; OLIGOMERIC SIZE; SERUM-ALBUMIN; GLYCATION

9/9,K/5 (Item 2 from file: 34)

DIALOG(R)File 34:SciSearch(R) Cited Ref Sci  
(c) 2007 The Thomson Corp. All rts. reserv.

14246175 Genuine Article#: 953HG Number of References: 44

**Title: Modulation of alpha- crystallin chaperone activity in diabetic rat lens by curcumin**

Author(s): Kumar PA; Suryanarayana P; Reddy PY; Reddy GB (REPRINT)

Corporate Source: Natl Inst Nutr,Hyderabad 500007/Andhra Pradesh/India/(REPRINT); Natl Inst Nutr,Hyderabad 500007/Andhra Pradesh/India/(geereddy@yahoo.com)

Journal: MOLECULAR VISION, 2005, V11, N66 (JUL 26), P561-568

ISSN: 1090-0535 Publication date: 20050726

Publisher: MOLECULAR VISION, C/O JEFF BOATRIGHT, LAB B, 5500 EMORY EYE CENTER, 1327 CLIFTON RD, N E, ATLANTA, GA 30322 USA

Language: English Document Type: ARTICLE

Geographic Location: India

Journal Subject Category: BIOCHEMISTRY & MOLECULAR BIOLOGY; OPHTHALMOLOGY

**Abstract:** Purpose: A decline in the chaperone-like activity of eye lens alpha- **crystallin** in diabetic conditions has been reported. In this study, we investigated whether curcumin, a dietary antioxidant, can manipulate the chaperone-like activity of alpha- **crystallin** in diabetic rat lens.

Methods: A group of rats received ip injection of streptozotocin (STZ; 35 mg/kg body weight in buffer) to induce hyperglycemia, while another group of rats received only buffer as vehicle and served as control. STZ-treated rats were assigned to 3 groups and fed either no

curcumin or 0.002% or 0.01% curcumin, respectively. Cataract progression due to hyperglycemia was monitored with a slit lamp biomicroscope. At the end of 8 weeks animals were sacrificed and lenses were collected. alpha H- and alpha L-crystallins from a set of pooled lenses in each group were isolated by gel filtration. Chaperone activity, hydrophobicity, and secondary and tertiary structure of alpha H- and alpha L-crystallins were assessed by light scattering/spectroscopic methods.

Results: A decrease in chaperone-like activity of alpha H- and alpha L-crystallins was observed in STZ-treated diabetic rats. The declined chaperone-like activity due to hyperglycemia was associated with reduced hydrophobicity and altered secondary and tertiary structure of alpha H- and alpha L-crystallins. Interestingly, alpha H- and alpha L-crystallins isolated from curcumin fed diabetic rat lenses had shown improved chaperone-like activity as compared to alpha H- and alpha L-crystallins from untreated diabetic rat lens. Feeding of curcumin prevented the alterations in hydrophobicity and structural changes due to STZ-induced hyperglycemia. Modulation of functional and structural properties by curcumin was found to be greater with the alpha L- **crystallin** than alpha H- **crystallin** . Loss of chaperone activity of alpha- **crystallin** , particularly alpha L- **crystallin** , in diabetic rat lens could be attributed at least partly to increased oxidative stress. Being an antioxidant, curcumin feeding has prevented the loss of alpha- **crystallin** chaperone activity and delayed the progression and maturation of diabetic cataract.

Conclusions: We demonstrate that curcumin, at the levels close to dietary consumption, prevented the loss of chaperone-like activity of alpha- **crystallin** vis-a-vis cataractogenesis due to diabetes in rat lens.

Identifiers--KeyWord Plus(R): A- **CRYSTALLIN** ; B- **CRYSTALLIN** ; **ARGININE HYDROCHLORIDE** ; IN-VIVO; CATARACT; AGGREGATION; PROTECT; STRESS; INDIA; RISK

Cited References:

- \*INT DIAB FED, 2001, DIAB ATL 2000
- BAI F, 2003, V278, P36876, J BIOL CHEM
- BHAT SP, 2003, V60, P205, PROG DRUG RES
- BOYLE DL, 2003, V3, P3, BMC OPHTHALMOL
- BRADY JP, 1997, V94, P884, P NATL ACAD SCI USA
- CHERIAN M, 1995, V212, P184, BIOCHEM BIOPH RES CO
- CHERIAN M, 1995, V208, P675, BIOCHEM BIOPH RES CO
- CLARK JI, 1996, V93, P15185, P NATL ACAD SCI USA
- DAS KP, 1995, V369, P321, FEBS LETT
- DERHAM BK, 2002, V364, P711, BIOCHEM J 3
- DERHAM BK, 1997, V328, P763, BIOCHEM J 3
- GANEVA E, 2001, V2, P205, CURR PROTEIN PEPT SC
- GROENEN PJTA, 1994, V225, P1, EUR J BIOCHEM
- HANSON SRA, 2000, V71, P195, EXP EYE RES
- HARDING J, 1991, CATARACT BIOICH EPID
- HOOK DWA, 1997, V247, P380, EUR J BIOCHEM
- HORWITZ J, 1998, V290, P365, METHOD ENZYMOL
- HORWITZ J, 1992, V11, P817, CURR EYE RES
- HORWITZ J, 2003, V76, P145, EXP EYE RES
- HUANG FY, 2000, V273, P197, BIOCHEM BIOPH RES CO
- KADOR PF, 1985, V28, P841, J MED CHEM

KUMAR MS, 2005, V280, P21726, J BIOL CHEM  
 KUMAR MS, 2004, V379, P273, BIOCHEM J 2  
 KYSELOVA Z, 2004, V18, P129, J DIABETES COMPLICAT  
 MARES JA, 2004, V62, P28, NUTR REV  
 NIRMALAN PK, 2004, V88, P989, BRIT J OPHTHALMOL  
 PRADEEPA R, 2002, V116, P121, INDIAN J MED RES  
 RAMAN B, 1997, V272, P23559, J BIOL CHEM  
 REDDY GB, 2000, V275, P4565, J BIOL CHEM  
 REDDY G, 2002, V8, P298, MOL VIS  
 REDDY GB, 2001, V282, P712, BIOCHEM BIOPH RES CO  
 REEVES PG, 1997, V127, P838, J NUTR  
 SHARMA KK, 1998, V273, P8965, J BIOL CHEM  
 SPECTOR A, 2000, V16, P193, J OCUL PHARMACOL TH  
 SRINIVAS V, 2003, V12, P1262, PROTEIN SCI  
 SRINIVAS V, 2005, V11, P249, MOL VIS  
 SURYANARAYANA P, 2003, V9, P223, MOL VIS  
 SURYANARAYANA P, 2005, V46, P2092, INVEST OPHTH VIS SCI  
 THAMPI P, 2002, V229, P113, MOL CELL BIOCHEM  
 THAMPI P, 2002, V43, P3265, INVEST OPHTH VIS SCI  
 THAMPI P, 2003, V42, P11857, BIOCHEMISTRY-US  
 UGHADE SN, 1998, V46, P221, INDIAN J OPHTHALMOL  
 VANBOEKEL MAM, 1992, V314, P1, FEBS LETT  
 YANG FS, 2005, V280, P5892, J BIOL CHEM

**Title: Modulation of alpha- crystallin chaperone activity in diabetic rat lens by curcumin**

**Abstract:** Purpose: A decline in the chaperone-like activity of eye lens alpha- **crystallin** in diabetic conditions has been reported. In this study, we investigated whether curcumin, a dietary antioxidant, can manipulate the chaperone-like activity of alpha- **crystallin** in diabetic rat lens.

**Methods:** A group of rats received ip injection of streptozotocin (STZ...

...functional and structural properties by curcumin was found to be greater with the alpha L- **crystallin** than alpha H- **crystallin** . Loss of chaperone activity of alpha- **crystallin** , particularly alpha L- **crystallin** , in diabetic rat lens could be attributed at least partly to increased oxidative stress. Being an antioxidant, curcumin feeding has prevented the loss of alpha- **crystallin** chaperone activity and delayed the progression and maturation of diabetic cataract.

**Conclusions:** We demonstrate that...

...the levels close to dietary consumption, prevented the loss of chaperone-like activity of alpha- **crystallin** vis-a-vis cataractogenesis due to diabetes in rat lens.

...Identifiers--A- **CRYSTALLIN** ; B- **CRYSTALLIN** ; **ARGININE HYDROCHLORIDE** ; IN-VIVO; CATARACT; AGGREGATION; PROTECT; STRESS; INDIA; RISK  
 ? t s9/9,k/11-20

9/9,K/11 (Item 8 from file: 34)  
 DIALOG(R)File 34:SciSearch(R) Cited Ref Sci  
 (c) 2007 The Thomson Corp. All rts. reserv.

05732997 Genuine Article#: WU248 Number of References: 41



Title: Modulation of endogenous antioxidant enzymes by nitric oxide in rat C-6 glial cells

Author(s): Dobashi K; Pahan K; Chahal A; Singh I (REPRINT)

Corporate Source: MED UNIV S CAROLINA, DEPT PEDIAT, DIV DEV NEUROGENET, 171 ASHLEY AVE/CHARLESTON//SC/29425 (REPRINT); MED UNIV S CAROLINA, DEPT PEDIAT, DIV DEV NEUROGENET/CHARLESTON//SC/29425

Journal: JOURNAL OF NEUROCHEMISTRY, 1997, V68, N5 (MAY), P1896-1903

ISSN: 0022-3042 Publication date: 19970500

Publisher: LIPPINCOTT-RAVEN PUBL, 227 EAST WASHINGTON SQ, PHILADELPHIA, PA 19106

Language: English Document Type: ARTICLE

Geographic Location: USA

Subfile: CC LIFE--Current Contents, Life Sciences

Journal Subject Category: BIOCHEMISTRY & MOLECULAR BIOLOGY; NEUROSCIENCES

Abstract: To understand the possible mechanism of nitric oxide

(NO)-mediated cytotoxicity, we investigated the effect of NO on the endogenous antioxidant enzymes (AOEs) catalase, glutathione peroxidase (GPX), and CuZn- and Mn-superoxide dismutases (SODs) in rat C-6 glial cells under conditions in which these cells expressed oligodendrocyte-like properties as evidenced by the expression of 2',3'-cyclic-nucleotide 3'-phosphohydrolase. The 24-h treatment with S-nitroso-N-acetylpenicillamine (SNAP), a NO donor, decreased the activities and the protein levels of catalase, GPX, and Mn-SOD in a dose-dependent manner. Alternatively, the activity and the protein level of CuZn-SOD were increased. 2-Phenyl-4,4,5,5-tetramethylimidazoline-1-oxyl-3-oxide (PTIO), a NO scavenger, blocked the effect of SNAP. Moreover, the treatment of C-6 cells with sodium nitroprusside, another NO donor, or with a combination of lipopolysaccharide (LPS) and interferon-gamma (IFN-gamma), which induce excessive production of NO, also significantly modulated the AOE activities in a manner similar to that seen with SNAP treatment. The compounds/enzymes that inhibit the production of NO (e.g., N-nitro-L-arginine methyl ester hydrochloride, arginase, and PTIO) blocked the effects of LPS and IFN-gamma on the activities of AOEs. Treatment with SNAP and a combination of LPS and IFN-gamma also modulated the mRNA levels of AOEs, parallel to the changes in their protein levels and activities, except for Mn-SOD where the combination of LPS and IFN-gamma markedly stimulated the mRNA expression. In spite of the stimulation of mRNA level, LPS and IFN-gamma significantly inhibited the activity of Mn-SOD within the first 24 h of incubation; however, Mn-SOD activity gradually increased with the increase in time of incubation. These results suggest that alterations in the status of AOEs by NO may be the basis of NO-induced cytotoxicity in disease states associated with excessive NO production.

Descriptors--Author Keywords: nitric oxide ; cytokine ; glia ; antioxidant enzymes ; gene expression

Identifiers--KeyWord Plus(R): SUPEROXIDE-DISMUTASE; REVERSIBLE BINDING; SYNTHASE ACTIVITY; INDUCTION; INHIBITION; MECHANISM; PEROXYNITRITE; CYTOTOXICITY; EXPRESSION; PROTEIN

Research Fronts: 95-2212 003 (PEROXYNITRITE IN-VITRO; NITRIC-OXIDE SYNTHASE; HYDROXYL RADICAL; FORMATION OF 8-NITROGUANINE; PC12 CELLS)

95-2984 001 (INDUCIBLE NITRIC-OXIDE SYNTHASE; CULTURED RAT ASTROCYTES; INCREASED EXPRESSION OF NADPH DIAPHORASE)

95-3190 001 (INCREASED ABUNDANCE OF SPECIFIC SKELETAL-MUSCLE PROTEIN-TYROSINE PHOSPHATASES; ALPHA-B-CRYSTALLIN EXPRESSION)

95-3891 001 (IN-VITRO TUMOR-NECROSIS-FACTOR CYTOTOXICITY; TRANSFECTION OF CELLS; MANGANESE SUPEROXIDE-DISMUTASE; GENE-EXPRESSION FOR

IMMUNOMODULATING CYTOKINES)

95-6776 001 (INDUCIBLE NITRIC-OXIDE SYNTHASE; RAT MACROPHAGES MEDIATE FUNGISTATIC ACTIVITY; MICROGLIAL RELEASE)

95-8090 001 (NITRIC-OXIDE SYNTHASE; INCREASED INTRACELLULAR CA<sup>2+</sup> SELECTIVELY SUPPRESSES IL-1-INDUCED NO PRODUCTION; HUMAN CENTRAL-NERVOUS-SYSTEM TUMORS)

Cited References:

ABERNETHY JL, 1974, V249, P7339, J BIOL CHEM  
AMES BN, 1996, V7, P115, METHOD ENZYMOL  
ANGGARD E, 1994, V343, P1196, LANCET  
ASAHI M, 1995, V270, P21035, J BIOL CHEM  
ASAYAMA K, 1986, V46, P604, J NEUROCHEM  
BAUDHUIN P, 1964, V92, P179, BIOCHEM J  
BECKMAN JS, 1990, V87, P1620, P NATL ACAD SCI USA  
BO L, 1994, V36, P778, ANN NEUROL  
BOLANOS JP, 1994, V63, P910, J NEUROCHEM  
BRADFORD MM, 1976, V72, P248, ANAL BIOCHEM  
BROWN GC, 1995, V232, P188, EUR J BIOCHEM  
BRUNE B, 1994, V54, P61, LIFE SCI  
CROSS AH, 1994, V93, P2684, J CLIN INVEST  
DORHEIM MA, 1994, V205, P659, BIOCHEM BIOPH RES CO  
DRUMMOND GI, 1971, V28, P153, BRAIN RES  
FEINSTEIN DL, 1994, V62, P315, J NEUROCHEM  
GALEA E, 1992, V89, P10945, P NATL ACAD SCI USA  
GHOSH B, 1996, V21, P533, FREE RACIC BIOL MED  
HENRY Y, 1993, V7, P1124, FASEB J  
ISCHIROPOULOS H, 1992, V298, P431, ARCH BIOCHEM BIOPHYS  
JAFFREY SR, 1995, V11, P417, ANN REV CELL BIOL  
KIFLE Y, 1996, V66, P2128, J NEUROCHEM  
KOPROWSKI H, 1993, V90, P3024, P NATL ACAD SCI USA  
KRONCKE KD, 1994, V200, P1105, BIOCHEM BIOPH RES CO  
KRONCKE KD, 1993, V1182, P221, BIOCHIM BIOPHYS ACTA  
LIPTON SA, 1993, V364, P626, NATURE  
MACKENZIEGRAHAM AJ, 1994, V16, P162, DEV NEUROSCI-BASEL  
MARLETTA MA, 1994, V78, P927, CELL  
MERRILL JE, 1993, V151, P2132, J IMMUNOL  
MITROVIC B, 1994, V61, P575, NEUROSCIENCE  
NATHAN C, 1994, V78, P915, CELL  
NORONHADUTRA AA, 1993, V321, P59, FEBS LETT  
OHSHIMA H, 1994, V305, P253, MUTAT RES  
PARK SK, 1994, V39, P405, J NEUROSCI RES  
RIBEIRO JMC, 1993, V260, P539, SCIENCE  
SIMMONS ML, 1992, V59, P897, J NEUROCHEM  
SPITZ DR, 1989, V179, P8, ANAL BIOCHEM  
STAMLER JS, 1994, V78, P913, CELL  
STUEHR DJ, 1989, V169, P1543, J EXP MED  
VOLPE JJ, 1983, V40, P530, J NEUROCHEM  
WONG GHW, 1988, V242, P941, SCIENCE

...Abstract: treatment. The compounds/enzymes that inhibit the production of NO (e.g., N-nitro-L- arginine methyl ester hydrochloride , arginase, and PTIO) blocked the effects of LPS and IFN-gamma on the activities of...

...Research Fronts: DIAPHORASE)

95-3190 001 (INCREASED ABUNDANCE OF SPECIFIC SKELETAL-MUSCLE PROTEIN-TYROSINE PHOSPHATASES; ALPHA-B- CRYSTALLIN EXPRESSION)

95-3891 001 (IN-VITRO TUMOR-NECROSIS-FACTOR CYTOTOXICITY; TRANSFECTION

OF CELLS; MANGANESE SUPEROXIDE...

9/9,K/12 (Item 1 from file: 35)  
DIALOG(R)File 35:Dissertation Abs Online  
(c) 2006 ProQuest Info&Learning. All rts. reserv.

02045668 ORDER NO: AADAA-INQ94247

Intragenic complementation and protein oligomerization studies in  
argininosuccinate lyase and its homologue delta crystallin

Author: Yu, Bomina

Degree: Ph.D.

Year: 2004

Corporate Source/Institution: University of Toronto (Canada) (0779)

Advisers: P. L. Honell; A. R. Davidson

Source: VOLUME 65/10-B OF DISSERTATION ABSTRACTS INTERNATIONAL.

PAGE 5140. 261 PAGES

Descriptors: CHEMISTRY, BIOCHEMISTRY

Descriptor Codes: 0487

ISBN: 0-612-94247-3

<?Pub Inc> Argininosuccinate lyase (ASL) is a ubiquitous enzyme that catalyzes the reversible cleavage of argininosuccinate to **arginine** and fumarate. This reaction is important both for the detoxification of ammonia *via* the urea cycle and for the biosynthesis of **arginine**. Through a process called 'gene sharing' ASL was recruited to the eye lens of birds and reptiles where it acts as the major lens **crystallin**. Both ASL and  $\delta$  **crystallin** exist as homotetramers with a monomer molecular weight of approximately 50 kDa. Extensive intragenic complementation was observed at the ASL locus in humans. Intragenic complementation occurs when certain combinations of mutant alleles produce an enzyme with greater catalytic activity than is observed in the homozygous state of either mutant.

In this thesis, ASL and  $\delta 2$  **crystallin** were used as model systems to study intragenic complementation and protein oligomerization. The structure and function of mutant proteins possessing amino acid substitutions associated with ASL deficiency were characterized. Mutations were found to either disturb the active site or drastically destabilize the protein such that proper metabolic function would be compromised. By coexpressing different pairs of mutants, intragenic complementation was found to occur between two active site mutants by the regeneration of native-like active sites and between stable and unstable mutants due to the increase in stability upon oligomerization. Complementation was also observed between  $\delta 2$  **crystallin** mutants and between  $\delta 2$  **crystallin** and ASL, implying that ASL and  $\delta 2$  **crystallin** have similar subunit interfaces and that the two proteins fold in the same manner. Both proteins were found to unfold in guanidine **hydrochloride** *via* a partially folded dimeric intermediate. Extensive site-directed mutagenesis at the subunit interface and of residues buried in the monomer suggested that unfolding occurs independent of the oligomeric state and illustrated the importance of inter-subunit salt-bridges for maintaining tetramer stability. Intragenic complementation was observed between mutant proteins with opposite amino acid substitutions in the subunit interface. Together these studies have provided insight into the pathology of argininosuccinic aciduria and the folding mechanism of ASL and  $\delta 2$  **crystallin**, and have illustrated the value of intragenic complementation studies when examining subunit interactions in oligomeric

proteins.

**Intragenic complementation and protein oligomerization studies in argininosuccinate lyase and its homologue delta crystallin**

...Argininosuccinate lyase (ASL) is a ubiquitous enzyme that catalyzes the reversible cleavage of argininosuccinate to **arginine** and fumarate. This reaction is important both for the detoxification of ammonia *<italic>via</italic>* the urea cycle and for the biosynthesis of **arginine**. Through a process called 'gene sharing' ASL was recruited to the eye lens of birds and reptiles where it acts as the major lens **crystallin**. Both ASL and  $\delta$  **crystallin** exist as homotetramers with a monomer molecular weight of approximately 50 kDa. Extensive intragenic complementation...

...observed in the homozygous state of either mutant.

In this thesis, ASL and  $\delta 2$  **crystallin** were used as model systems to study intragenic complementation and protein oligomerization. The structure and...

...due to the increase in stability upon oligomerization. Complementation was also observed between  $\delta 2$  **crystallin** mutants and between  $\delta 2$  **crystallin** and ASL, implying that ASL and  $\delta 2$  **crystallin** have similar subunit interfaces and that the two proteins fold in the same manner. Both proteins were found to unfold in guanidine **hydrochloride** *<italic>via</italic>* a partially folded dimeric intermediate. Extensive site-directed mutagenesis at the subunit...

...into the pathology of argininosuccinic aciduria and the folding mechanism of ASL and  $\delta 2$  **crystallin**, and have illustrated the value of intragenic complementation studies when examining subunit interactions in oligomeric...

9/9,K/13 (Item 1 from file: 71)

DIALOG(R)File 71:ELSEVIER BIOBASE

(c) 2007 Elsevier B.V. All rts. reserv.

03320778

2006097921

**Effect of site-directed mutagenesis of methylglyoxal-modifiable arginine residues on the structure and chaperone function of human alphaA-crystallin**

Biswas A.; Miller A.; Oya-Ito T.; Santhoshkumar P.; Bhat M.; Nagaraj R.H.  
ADDRESS: R.H. Nagaraj, Department of Ophthalmology, Case Western Reserve University, Cleveland, OH 44106, United States

EMAIL: ram.nagaraj@case.edu

Journal: Biochemistry, 45/14 (4569-4577), 2006, United States

PUBLICATION DATE: April 11, 2006

CODEN: BICHA

ISSN: 0006-2960

DOCUMENT TYPE: Article

LANGUAGES: English

SUMMARY LANGUAGES: English

NO. OF REFERENCES: 59

We reported previously that chemical modification of human alphaA-**crystallin** by a metabolic dicarbonyl compound, methylglyoxal (MGO), enhances its chaperone-like function, a phenomenon which we attributed to formation of argpyrimidine at **arginine** residues (R) 21, 49, and 103. This

structural change removes the positive charge on the **arginine** residues. To explore this mechanism further, we replaced these three R residues with a neutral alanine (A) residue one at a time or in combination and examined the impact on the structure and chaperone function. Measurement of intrinsic tryptophan fluorescence and near-UV CD spectra revealed alteration of the microenvironment of aromatic amino acid residues in mutant proteins. When compared to wild-type (wt) **alphaA-crystallin**, the chaperone function of R21A and R103A mutants increased 20% and 18% as measured by the insulin aggregation assay and increased it as much as 39% and 28% when measured by the citrate synthase (CS) aggregation assay. While the R49A mutant lost most of its chaperone function, R21A/R103A and R21A/R49A/R103A mutants had slightly better function (6-14% and 10-14%) than the wt protein in these assays. R21A and R103A mutants had higher surface hydrophobicity than wt **alphaA-crystallin**, but the R49A mutant had lower hydrophobicity. R21A and R103A mutants, but not the R49A mutant, were more efficient than wt protein in refolding guanidine **hydrochloride**-treated malate dehydrogenase to its native state. Our findings indicate that the positive charges on R21, R49, and R103 are important determinants of the chaperone function of **alphaA-crystallin** and suggest that chemical modification of **arginine** residues may play a role in protein aggregation during lens aging and cataract formation. (c) 2006 American Chemical Society.

CLASSIFICATION CODE AND DESCRIPTION:

- 82.3.6 - PROTEIN BIOCHEMISTRY / PROTEIN ENGINEERING / Mutation, Expression and Isolation
- 82.2.8 - PROTEIN BIOCHEMISTRY / STRUCTURAL STUDIES / Folding, Unfolding and Stability
- 82.2.3 - PROTEIN BIOCHEMISTRY / STRUCTURAL STUDIES / Protein Crystallization and Crystal Structures

**Effect of site-directed mutagenesis of methylglyoxal-modifiable arginine residues on the structure and chaperone function of human alphaA-crystallin**

We reported previously that chemical modification of human **alphaA-crystallin** by a metabolic dicarbonyl compound, methylglyoxal (MGO), enhances its chaperone-like function, a phenomenon which we attributed to formation of argpyrimidine at **arginine** residues (R) 21, 49, and 103. This structural change removes the positive charge on the **arginine** residues. To explore this mechanism further, we replaced these three R residues with a neutral...

...of aromatic amino acid residues in mutant proteins. When compared to wild-type (wt) **alphaA-crystallin**, the chaperone function of R21A and R103A mutants increased 20% and 18% as measured by...

...protein in these assays. R21A and R103A mutants had higher surface hydrophobicity than wt **alphaA-crystallin**, but the R49A mutant had lower hydrophobicity. R21A and R103A mutants, but not the R49A mutant, were more efficient than wt protein in refolding guanidine **hydrochloride**-treated malate dehydrogenase to its native state. Our findings indicate that the positive charges on R21, R49, and R103 are important determinants of the chaperone function of **alphaA-crystallin** and suggest that chemical modification of **arginine** residues may play a role in protein aggregation during lens aging and cataract formation. (c...

9/9,K/14 (Item 2 from file: 71)  
DIALOG(R) File 71:ELSEVIER BIOBASE  
(c) 2007 Elsevier B.V. All rts. reserv.

02344455 2003128140

**Structural perturbation and enhancement of the chaperone-like activity of  
alpha- crystallin by arginine hydrochloride**

Srinivas V.; Raman B.; Rao K.S.; Ramakrishna T.; Rao Ch.M.

ADDRESS: Ch.M. Rao, Ctr. for Cell. and Molecular Biology, Uppal Road,  
Hyderabad 500 007, India

EMAIL: mohan@ccmb.res.in

Journal: Protein Science, 12/6 (1262-1270), 2003, United States

PUBLICATION DATE: June 1, 2003

CODEN: PRCIE

ISSN: 0961-8368

DOCUMENT TYPE: Article

LANGUAGES: English SUMMARY LANGUAGES: English

NO. OF REFERENCES: 57

Structural perturbation of alpha- **crystallin** is shown to enhance its molecular chaperone-like activity in preventing aggregation of target proteins. We demonstrate that **arginine**, a biologically compatible molecule that is known to bind to the peptide backbone and negatively charged side-chains, increases the chaperone-like activity of calf eye lens alpha- **crystallin** as well as recombinant human alphaA- and alphaB-crystallins. Arginine-induced increase in the chaperone activity is more pronounced for alphaB- **crystallin** than for alphaA- **crystallin**. Other guanidinium compounds such as aminoguanidine **hydrochloride** and guanidine **hydrochloride** also show a similar effect, but to different extents. A point mutation, R120G, in alphaB- **crystallin** that is associated with desmin-related myopathy, results in a significant loss of chaperone-like activity. **Arginine** restores the activity of mutant protein to a considerable extent. We have investigated the effect of **arginine** on the structural changes of alpha- **crystallin** by circular dichroism, fluorescence, and glycerol gradient sedimentation. Far-UV CD spectra show no significant changes in secondary structure, whereas near-UV CD spectra show subtle changes in the presence of **arginine**. Glycerol gradient sedimentation shows a significant decrease in the size of alpha- **crystallin** oligomer in the presence of **arginine**. Increased exposure of hydrophobic surfaces of alpha- **crystallin**, as monitored by pyrene-solubilization and ANS-fluorescence, is observed in the presence of **arginine**. These results show that **arginine** brings about subtle changes in the tertiary structure and significant changes in the quaternary structure of alpha- **crystallin** and enhances its chaperone-like activity significantly. This study should prove useful in designing strategies to improve chaperone function for therapeutic applications.

DESCRIPTORS:

Chaperone-like activity; alpha- **crystallin**; **Arginine**; Aminoguanidine;  
Structural perturbation

CLASSIFICATION CODE AND DESCRIPTION:

82.2.8 - PROTEIN BIOCHEMISTRY / STRUCTURAL STUDIES / Folding, Unfolding and  
Stability

**Structural perturbation and enhancement of the chaperone-like activity of  
alpha- crystallin by arginine hydrochloride**

Structural perturbation of **alpha-crystallin** is shown to enhance its molecular chaperone-like activity in preventing aggregation of target proteins. We demonstrate that **arginine**, a biologically compatible molecule that is known to bind to the peptide backbone and negatively charged side-chains, increases the chaperone-like activity of calf eye lens **alpha-crystallin** as well as recombinant human **alphaA-** and **alphaB-crystallins**. Arginine-induced increase in the chaperone activity is more pronounced for **alphaB-crystallin** than for **alphaA-crystallin**. Other guanidinium compounds such as aminoguanidine **hydrochloride** and guanidine **hydrochloride** also show a similar effect, but to different extents. A point mutation, R120G, in **alphaB-crystallin** that is associated with desmin-related myopathy, results in a significant loss of chaperone-like activity. **Arginine** restores the activity of mutant protein to a considerable extent. We have investigated the effect of **arginine** on the structural changes of **alpha-crystallin** by circular dichroism, fluorescence, and glycerol gradient sedimentation. Far-UV CD spectra show no significant...

...in secondary structure, whereas near-UV CD spectra show subtle changes in the presence of **arginine**. Glycerol gradient sedimentation shows a significant decrease in the size of **alpha-crystallin** oligomer in the presence of **arginine**. Increased exposure of hydrophobic surfaces of **alpha-crystallin**, as monitored by pyrene-solubilization and ANS-fluorescence, is observed in the presence of **arginine**. These results show that **arginine** brings about subtle changes in the tertiary structure and significant changes in the quaternary structure of **alpha-crystallin** and enhances its chaperone-like activity significantly. This study should prove useful in designing strategies...

DESCRIPTORS:

Chaperone-like activity; **alpha-crystallin**; **Arginine**; Aminoguanidine; Structural perturbation

9/9,K/15 (Item 1 from file: 73)

DIALOG(R) File 73:EMBASE

(c) 2007 Elsevier B.V. All rts. reserv.

13754273 EMBASE No: 2006169408

**Effect of site-directed mutagenesis of methylglyoxal-modifiable arginine residues on the structure and chaperone function of human alphaA-crystallin**

Biswas A.; Miller A.; Oya-Ito T.; Santhoshkumar P.; Bhat M.; Nagaraj R.H. R.H. Nagaraj, Department of Ophthalmology, Case Western Reserve University, Cleveland, OH 44106 United States

AUTHOR EMAIL: ram.nagaraj@case.edu

Biochemistry ( BIOCHEMISTRY ) (United States) 11 APR 2006, 45/14 (4569-4577)

CODEN: BICHA ISSN: 0006-2960

DOCUMENT TYPE: Journal ; Article

LANGUAGE: ENGLISH SUMMARY LANGUAGE: ENGLISH

NUMBER OF REFERENCES: 59

We reported previously that chemical modification of human **alphaA-crystallin** by a metabolic dicarbonyl compound, methylglyoxal (MGO), enhances its chaperone-like function, a phenomenon which we attributed to formation of argpyrimidine at **arginine** residues (R) 21, 49, and 103. This

structural change removes the positive charge on the **arginine** residues. To explore this mechanism further, we replaced these three R residues with a neutral alanine (A) residue one at a time or in combination and examined the impact on the structure and chaperone function. Measurement of intrinsic tryptophan fluorescence and near-UV CD spectra revealed alteration of the microenvironment of aromatic amino acid residues in mutant proteins. When compared to wild-type (wt) **alphaA-crystallin**, the chaperone function of R21A and R103A mutants increased 20% and 18% as measured by the insulin aggregation assay and increased it as much as 39% and 28% when measured by the citrate synthase (CS) aggregation assay. While the R49A mutant lost most of its chaperone function, R21A/R103A and R21A/R49A/R103A mutants had slightly better function (6-14% and 10-14%) than the wt protein in these assays. R21A and R103A mutants had higher surface hydrophobicity than wt **alphaA-crystallin**, but the R49A mutant had lower hydrophobicity. R21A and R103A mutants, but not the R49A mutant, were more efficient than wt protein in refolding guanidine **hydrochloride**-treated malate dehydrogenase to its native state. Our findings indicate that the positive charges on R21, R49, and R103 are important determinants of the chaperone function of **alphaA-crystallin** and suggest that chemical modification of **arginine** residues may play a role in protein aggregation during lens aging and cataract formation. (c) 2006 American Chemical Society.

#### DRUG DESCRIPTORS:

\***alpha crystallin**; \*methylglyoxal; \* **arginine**  
chaperone; pyrimidine; amino acid; citrate synthase; mutant protein;  
guanidine **hydrochloride**; insulin; malate dehydrogenase; aromatic amino acid

#### MEDICAL DESCRIPTORS:

\*protein structure; \*site directed mutagenesis; \*protein function  
protein modification; circular dichroism; amino acid substitution;  
hydrophobicity; protein folding; cataractogenesis; protein aggregation;  
aging; lens; human; controlled study; article; priority journal

CAS REGISTRY NO.: 78-98-8 (methylglyoxal); 1119-34-2, 15595-35-4, 7004-12-8  
, 74-79-3 ( **arginine** ); 289-95-2 (pyrimidine); 65072-01-7 (amino acid);  
9027-96-7 (citrate synthase); 50-01-1 (guanidine **hydrochloride** );  
9004-10-8 (insulin); 9001-64-3 (malate dehydrogenase)

#### SECTION HEADINGS:

012 Ophthalmology  
029 Clinical and Experimental Biochemistry

**Effect of site-directed mutagenesis of methylglyoxal-modifiable arginine residues on the structure and chaperone function of human alphaA-crystallin**

We reported previously that chemical modification of human **alphaA-crystallin** by a metabolic dicarbonyl compound, methylglyoxal (MGO), enhances its chaperone-like function, a phenomenon which we attributed to formation of argpyrimidine at **arginine** residues (R) 21, 49, and 103. This structural change removes the positive charge on the **arginine** residues. To explore this mechanism further, we replaced these three R residues with a neutral...

...of aromatic amino acid residues in mutant proteins. When compared to wild-type (wt) **alphaA-crystallin**, the chaperone function of R21A and R103A mutants increased 20% and 18% as measured by...



...protein in these assays. R21A and R103A mutants had higher surface hydrophobicity than wt alphaA- **crystallin** , but the R49A mutant had lower hydrophobicity. R21A and R103A mutants, but not the R49A mutant, were more efficient than wt protein in refolding guanidine **hydrochloride** -treated malate dehydrogenase to its native state. Our findings indicate that the positive charges on R21, R49, and R103 are important determinants of the chaperone function of alphaA- **crystallin** and suggest that chemical modification of **arginine** residues may play a role in protein aggregation during lens aging and cataract formation. (c...

DRUG DESCRIPTORS:

\*alpha **crystallin** ; \*methylglyoxal; \* **arginine**  
chaperone; pyrimidine; amino acid; citrate synthase; mutant protein;  
guanidine **hydrochloride** ; insulin; malate dehydrogenase; aromatic amino acid

...CAS REGISTRY NO.: 74-79-3 ( **arginine** ); 289-95-2 (pyrimidine);  
65072-01-7 (amino acid); 9027-96-7 (citrate synthase); 50-01-1 ( **guanidine hydrochloride** ); 9004-10-8 (insulin); 9001-64-3 (malate dehydrogenase)

9/9,K/16 (Item 2 from file: 73)

DIALOG(R)File 73:EMBASE

(c) 2007 Elsevier B.V. All rts. reserv.

13357267 EMBASE No: 2005431360

**Arginine hydrochloride enhances the dynamics of subunit assembly and the chaperone-like activity of alpha- crystallin**

Srinivas V.; Raman B.; Sridhar Rao K.; Ramakrishna T.; Mohan Rao Ch.  
Dr. Ch. Mohan Rao, Centre for Cellular and Molecular Biology, Uppal Road,  
Hyderabad 500 007 India

AUTHOR EMAIL: mohan@ccmb.res.in

Molecular Vision ( MOL. VISION ) (United States) 01 APR 2005, 11/-  
(249-255)

CODEN: MVEPF ISSN: 1090-0535

DOCUMENT TYPE: Journal ; Article

LANGUAGE: ENGLISH SUMMARY LANGUAGE: ENGLISH

NUMBER OF REFERENCES: 49

Purpose: alpha- **Crystallin** , a major eye lens protein, bears homology with small heat shock proteins (sHsps) and exhibits molecular chaperone-like activity. Structural perturbation by temperature or low concentrations of denaturants leads to enhancement of its chaperone-like activity. We have earlier demonstrated similar enhancement of chaperone-like activity using biologically compatible solutes such as **arginine hydrochloride** and aminoguanidine. The purpose of the present study is to get an insight into the mechanism of the **arginine** induced enhancement of chaperone-like activity of **crystallin** . Methods: The effect of **arginine hydrochloride** on the chaperone-like activity of alpha-**crystallin** at 25 degreesC was studied using DTT induced aggregation of insulin as a model system. Changes in the accessibility of the thiol group near the end of the a- **crystallin** domain in the absence and the presence of **arginine hydrochloride** were studied using dithiobisnitrobenzoic acid. Fluorescence resonance energy transfer studies were performed to investigate changes in the dynamics of the subunit assembly. Urea induced denaturation studies of alpha- **crystallin** were carried out to investigate structural destabilization of alpha- **crystallin** , if any, in the presence of **arginine hydrochloride** . Results: **Arginine hydrochloride**

increases the chaperone-like activity of alpha- **crystallin** several fold towards DTT induced aggregation of insulin at room temperature. Our study shows that both the extent and the rate of accessibility of the thiol group are increased in the presence of **arginine** . Fluorescence resonance energy transfer experiments show that **arginine hydrochloride** significantly increases the subunit exchange between the oligomers of alpha- **crystallin** .

**Arginine** induced structural perturbation and loosening of subunit assembly of alpha- **crystallin** leads to overall destabilization of the protein as reflected by the urea denaturation study. Conclusions: **Arginine** perturbs the tertiary and quaternary structure of a- **crystallin** and enhances the dynamics of the subunit assembly leading to enhanced chaperone-like activity. Thus, in addition to size, surface hydrophobicity, and charge distribution, the dynamics of the subunit assembly appears to be one of the critical factors that can modulate the chaperone activity.

(c)2005 Molecular Vision.

#### DRUG DESCRIPTORS:

\* **arginine** ; \*chaperone; \*alpha **crystallin**  
dithiothreitol; insulin; benzoic acid; urea

#### MEDICAL DESCRIPTORS:

protein assembly; biological model; protein domain; fluorescence resonance energy transfer; protein denaturation; protein structure; room temperature; hydrophobicity; article; priority journal

CAS REGISTRY NO.: 1119-34-2, 15595-35-4, 7004-12-8, 74-79-3 ( **arginine** );  
3483-12-3 (dithiothreitol); 9004-10-8 (insulin); 532-32-1, 582-25-2,  
65-85-0, 766-76-7 (benzoic acid); 57-13-6 (urea)

#### SECTION HEADINGS:

029 Clinical and Experimental Biochemistry

**Arginine hydrochloride enhances the dynamics of subunit assembly and the chaperone-like activity of alpha- crystallin**

Purpose: alpha- **Crystallin** , a major eye lens protein, bears homology with small heat shock proteins (sHsps) and exhibits...

...have earlier demonstrated similar enhancement of chaperone-like activity using biologically compatible solutes such as **arginine hydrochloride** and aminoguanidine. The purpose of the present study is to get an insight into the mechanism of the **arginine** induced enhancement of chaperone-like activity of **crystallin** . Methods: The effect of **arginine hydrochloride** on the chaperone-like activity of alpha- **crystallin** at 25 degreesC was studied using DTT induced aggregation of insulin as a model system. Changes in the accessibility of the thiol group near the end of the a- **crystallin** domain in the absence and the presence of **arginine hydrochloride** were studied using dithiobisnitrobenzoic acid. Fluorescence resonance energy transfer studies were performed to investigate changes in the dynamics of the subunit assembly. Urea induced denaturation studies of alpha- **crystallin** were carried out to investigate structural destabilization of alpha- **crystallin** , if any, in the presence of **arginine hydrochloride** . Results: **Arginine hydrochloride** increases the chaperone-like activity of alpha- **crystallin** several fold towards DTT induced aggregation of insulin at room temperature. Our study shows that...

...and the rate of accessibility of the thiol group are increased in the presence of **arginine** . Fluorescence resonance energy transfer experiments show that **arginine hydrochloride** significantly increases the subunit exchange between the oligomers of alpha- **crystallin** . **Arginine** induced structural perturbation and loosening of subunit assembly of alpha-

**crystallin** leads to overall destabilization of the protein as reflected by the urea denaturation study. Conclusions: **Arginine** perturbs the tertiary and quaternary structure of **alpha-crystallin** and enhances the dynamics of the subunit assembly leading to enhanced chaperone-like activity. Thus...

DRUG DESCRIPTORS:

\* **arginine** ; \*chaperone; \***alpha crystallin**

...CAS REGISTRY NO.: 74-79-3 ( **arginine** ); 3483-12-3 (dithiothreitol);  
9004-10-8 (insulin); 532-32-1...

9/9,K/17 (Item 3 from file: 73)

DIALOG(R) File 73:EMBASE

(c) 2007 Elsevier B.V. All rts. reserv.

12095641 EMBASE No: 2003207170

**Structural perturbation and enhancement of the chaperone-like activity of alpha-crystallin by arginine hydrochloride**

Srinivas V.; Raman B.; Rao K.S.; Ramakrishna T.; Rao Ch.M.

Ch.M. Rao, Ctr. for Cell. and Molecular Biology, Uppal Road, Hyderabad  
500 007 India

AUTHOR EMAIL: mohan@ccmb.res.in

Protein Science ( PROTEIN SCI. ) (United States) 01 JUN 2003, 12/6  
(1262-1270)

CODEN: PRCIE ISSN: 0961-8368

DOCUMENT TYPE: Journal ; Article

LANGUAGE: ENGLISH SUMMARY LANGUAGE: ENGLISH

NUMBER OF REFERENCES: 57

Structural perturbation of **alpha-crystallin** is shown to enhance its molecular chaperone-like activity in preventing aggregation of target proteins. We demonstrate that **arginine**, a biologically compatible molecule that is known to bind to the peptide backbone and negatively charged side-chains, increases the chaperone-like activity of calf eye lens **alpha-crystallin** as well as recombinant human **alphaA-** and **alphaB-crystallins**. Arginine-induced increase in the chaperone activity is more pronounced for **alphaB-crystallin** than for **alphaA-crystallin**. Other guanidinium compounds such as aminoguanidine **hydrochloride** and guanidine **hydrochloride** also show a similar effect, but to different extents. A point mutation, R120G, in **alphaB-crystallin** that is associated with desmin-related myopathy, results in a significant loss of chaperone-like activity. **Arginine** restores the activity of mutant protein to a considerable extent. We have investigated the effect of **arginine** on the structural changes of **alpha-crystallin** by circular dichroism, fluorescence, and glycerol gradient sedimentation. Far-UV CD spectra show no significant changes in secondary structure, whereas near-UV CD spectra show subtle changes in the presence of **arginine**. Glycerol gradient sedimentation shows a significant decrease in the size of **alpha-crystallin** oligomer in the presence of **arginine**. Increased exposure of hydrophobic surfaces of **alpha-crystallin**, as monitored by pyrene-solubilization and ANS-fluorescence, is observed in the presence of **arginine**. These results show that **arginine** brings about subtle changes in the tertiary structure and significant changes in the quaternary structure of **alpha-crystallin** and enhances its chaperone-like activity significantly. This study should prove useful in designing strategies to improve chaperone function for therapeutic applications.

DRUG DESCRIPTORS:

\*chaperone; \*alpha **crystallin** --endogenous compound--ec; \* **arginine**  
guanidine derivative; aminoguanidine; guanidine **hydrochloride** ; desmin;  
glycerol; pyrene; oligomer; unclassified drug

MEDICAL DESCRIPTORS:

\*protein structure

structure analysis; protein targeting; protein binding; point mutation;  
circular dichroism; fluorescence; sedimentation; protein secondary  
structure; solubilization; protein tertiary structure; protein quaternary  
structure; nonhuman; article; priority journal

DRUG TERMS (UNCONTROLLED): alpha b **crystallin**

CAS REGISTRY NO.: 1119-34-2, 15595-35-4, 7004-12-8, 74-79-3 ( **arginine** );  
1068-42-4, 2582-30-1, 79-17-4 (aminoguanidine); 50-01-1 (guanidine  
**hydrochloride** ); 56-81-5 (glycerol); 129-00-0 (pyrene)

SECTION HEADINGS:

029 Clinical and Experimental Biochemistry

**Structural perturbation and enhancement of the chaperone-like activity of  
alpha- crystallin by arginine hydrochloride**

Structural perturbation of alpha- **crystallin** is shown to enhance its  
molecular chaperone-like activity in preventing aggregation of target  
proteins. We demonstrate that **arginine** , a biologically compatible  
molecule that is known to bind to the peptide backbone and negatively  
charged side-chains, increases the chaperone-like activity of calf eye lens  
alpha- **crystallin** as well as recombinant human alphaA- and  
alphaB-crystallins. Arginine-induced increase in the chaperone activity is  
more pronounced for alphaB- **crystallin** than for alphaA- **crystallin** . Other  
guanidinium compounds such as aminoguanidine **hydrochloride** and guanidine  
**hydrochloride** also show a similar effect, but to different extents. A  
point mutation, R120G, in alphaB- **crystallin** that is associated with  
desmin-related myopathy, results in a significant loss of chaperone-like  
activity. **Arginine** restores the activity of mutant protein to a  
considerable extent. We have investigated the effect of **arginine** on the  
structural changes of alpha- **crystallin** by circular dichroism,  
fluorescence, and glycerol gradient sedimentation. Far-UV CD spectra show  
no significant...

...in secondary structure, whereas near-UV CD spectra show subtle changes  
in the presence of **arginine** . Glycerol gradient sedimentation shows a  
significant decrease in the size of alpha- **crystallin** oligomer in the  
presence of **arginine** . Increased exposure of hydrophobic surfaces of  
alpha- **crystallin** , as monitored by pyrene-solubilization and  
ANS-fluorescence, is observed in the presence of **arginine** . These results  
show that **arginine** brings about subtle changes in the tertiary structure  
and significant changes in the quaternary structure of alpha- **crystallin**  
and enhances its chaperone-like activity significantly. This study should  
prove useful in designing strategies...

DRUG DESCRIPTORS:

\*chaperone; \*alpha **crystallin** --endogenous compound--ec; \* **arginine**  
guanidine derivative; aminoguanidine; guanidine **hydrochloride** ; desmin;  
glycerol; pyrene; oligomer; unclassified drug

DRUG TERMS (UNCONTROLLED): alpha b **crystallin**

...CAS REGISTRY NO.: 74-79-3 ( **arginine** ); 1068-42-4...

...79-17-4 (aminoguanidine); 50-01-1 (guanidine **hydrochloride** ); 56-81-5 (  
glycerol); 129-00-0 (pyrene)

9/9,K/18 (Item 1 from file: 155)  
DIALOG(R)File 155:MEDLINE(R)  
(c) format only 2006 Dialog. All rts. reserv.

20818450 PMID: 16584192

**Effect of site-directed mutagenesis of methylglyoxal-modifiable arginine residues on the structure and chaperone function of human alphaA-crystallin .**

Biswas Ashis; Miller Antonia; Oya-Ito Tomoko; Santhoshkumar Puttur; Bhat Manjunatha; Nagaraj Ram H

Department of Ophthalmology, Case Western Reserve University, Cleveland, Ohio 44106, USA.

Biochemistry (United States) Apr 11 2006, 45 (14) p4569-77, ISSN 0006-2960--Print Journal Code: 0370623

Contract/Grant No.: P30EY-11373; EY; NEI; R01EY-016219; EY; NEI; R01EY-09912; EY; NEI

Publishing Model Print

Document type: Journal Article

Languages: ENGLISH

Main Citation Owner: NLM

Record type: MEDLINE; Completed

Subfile: INDEX MEDICUS; Toxbib

We reported previously that chemical modification of human alphaA-crystallin by a metabolic dicarbonyl compound, methylglyoxal (MGO), enhances its chaperone-like function, a phenomenon which we attributed to formation of argpyrimidine at arginine residues (R) 21, 49, and 103. This structural change removes the positive charge on the arginine residues. To explore this mechanism further, we replaced these three R residues with a neutral alanine (A) residue one at a time or in combination and examined the impact on the structure and chaperone function. Measurement of intrinsic tryptophan fluorescence and near-UV CD spectra revealed alteration of the microenvironment of aromatic amino acid residues in mutant proteins. When compared to wild-type (wt) alphaA-crystallin, the chaperone function of R21A and R103A mutants increased 20% and 18% as measured by the insulin aggregation assay and increased it as much as 39% and 28% when measured by the citrate synthase (CS) aggregation assay. While the R49A mutant lost most of its chaperone function, R21A/R103A and R21A/R49A/R103A mutants had slightly better function (6-14% and 10-14%) than the wt protein in these assays. R21A and R103A mutants had higher surface hydrophobicity than wt alphaA-crystallin, but the R49A mutant had lower hydrophobicity. R21A and R103A mutants, but not the R49A mutant, were more efficient than wt protein in refolding guanidine hydrochloride-treated malate dehydrogenase to its native state. Our findings indicate that the positive charges on R21, R49, and R103 are important determinants of the chaperone function of alphaA-crystallin and suggest that chemical modification of arginine residues may play a role in protein aggregation during lens aging and cataract formation.

Descriptors: \*Arginine --physiology--PH; \*Molecular Chaperones --physiology--PH; \*alpha-Crystallin A Chain--physiology--PH; Arginine --chemistry--CH; Carbonic Anhydrases--metabolism--ME; Circular Dichroism; Humans; Mutagenesis, Site-Directed; Protein Structure, Secondary; Pyruvaldehyde--pharmacology--PD; Research Support, N.I.H., Extramural; Research Support, Non-U.S. Gov't; Spectrometry, Fluorescence; alpha-Crystallin A Chain--chemistry--CH

CAS Registry No.: 0 (Molecular Chaperones); 0 (alpha-Crystallin A Chain); 74-79-3 (Arginine); 78-98-8 (Pyruvaldehyde)

Enzyme No.: EC 4.2.1.1 (Carbonic Anhydrases)  
Record Date Created: 20060404  
Record Date Completed: 20060530

**Effect of site-directed mutagenesis of methylglyoxal-modifiable arginine residues on the structure and chaperone function of human alphaA-crystallin .**

We reported previously that chemical modification of human alphaA-crystallin by a metabolic dicarbonyl compound, methylglyoxal (MGO), enhances its chaperone-like function, a phenomenon which we attributed to formation of argpyrimidine at **arginine** residues (R) 21, 49, and 103. This structural change removes the positive charge on the **arginine** residues. To explore this mechanism further, we replaced these three R residues with a neutral...

... of aromatic amino acid residues in mutant proteins. When compared to wild-type (wt) alphaA- **crystallin** , the chaperone function of R21A and R103A mutants increased 20% and 18% as measured by...

... protein in these assays. R21A and R103A mutants had higher surface hydrophobicity than wt alphaA- **crystallin** , but the R49A mutant had lower hydrophobicity. R21A and R103A mutants, but not the R49A mutant, were more efficient than wt protein in refolding guanidine **hydrochloride** -treated malate dehydrogenase to its native state. Our findings indicate that the positive charges on R21, R49, and R103 are important determinants of the chaperone function of alphaA- **crystallin** and suggest that chemical modification of **arginine** residues may play a role in protein aggregation during lens aging and cataract formation.

Descriptors: **\*Arginine** --physiology--PH; **\*Molecular Chaperones** --physiology--PH; **\*alpha- Crystallin A Chain**--physiology--PH; **Arginine** --chemistry--CH; Carbonic Anhydrases--metabolism--ME; Circular Dichroism; Humans; Mutagenesis, Site-Directed; Protein Structure, Secondary...

...Support, N.I.H., Extramural; Research Support, Non-U.S. Gov't; Spectrometry, Fluorescence; alpha- **Crystallin A Chain**--chemistry--CH  
Chemical Name: Molecular Chaperones; alpha- **Crystallin A Chain**; **Arginine** ; Pyruvaldehyde; Carbonic Anhydrases

9/9,K/19 (Item 2 from file: 155)  
DIALOG(R) File 155:MEDLINE(R)  
(c) format only 2006 Dialog. All rts. reserv.

20362892 PMID: 15827547

**Arginine hydrochloride enhances the dynamics of subunit assembly and the chaperone-like activity of alpha- crystallin .**

Srinivas V; Raman B; Rao K Sridhar; Ramakrishna T; Rao Ch Mohan  
Centre for Cellular and Molecular Biology, Hyderabad, India.

Molecular vision electronic resource (United States) 2005, 11

p249-55, ISSN 1090-0535--Electronic Journal Code: 9605351

Publishing Model Electronic

Document type: Journal Article

Languages: ENGLISH

Main Citation Owner: NLM

Record type: MEDLINE; Completed

Subfile: INDEX MEDICUS

PURPOSE: Alpha- **crystallin** , a major eye lens protein, bears homology

with small heat shock proteins (sHsps) and exhibits molecular chaperone-like activity. Structural perturbation by temperature or low concentrations of denaturants leads to enhancement of its chaperone-like activity. We have earlier demonstrated similar enhancement of chaperone-like activity using biologically compatible solutes such as **arginine hydrochloride** and aminoguanidine. The purpose of the present study is to get an insight into the mechanism of the **arginine** induced enhancement of chaperone-like activity of **alpha-crystallin**. METHODS: The effect of **arginine hydrochloride** on the chaperone-like activity of **alpha-crystallin** at 25 degrees C was studied using DTT induced aggregation of insulin as a model system. Changes in the accessibility of the thiol group near the end of the **alpha-crystallin** domain in the absence and the presence of **arginine hydrochloride** were studied using dithiobisnitrobenzoic acid. Fluorescence resonance energy transfer studies were performed to investigate changes in the dynamics of the subunit assembly. Urea induced denaturation studies of **alpha-crystallin** were carried out to investigate structural destabilization of **alpha-crystallin**, if any, in the presence of **arginine hydrochloride**. RESULTS: **Arginine hydrochloride** increases the chaperone-like activity of **alpha-crystallin** several fold towards DTT induced aggregation of insulin at room temperature. Our study shows that both the extent and the rate of accessibility of the thiol group are increased in the presence of **arginine**. Fluorescence resonance energy transfer experiments show that **arginine hydrochloride** significantly increases the subunit exchange between the oligomers of **alpha-crystallin**. **Arginine** induced structural perturbation and loosening of subunit assembly of **alpha-crystallin** leads to overall destabilization of the protein as reflected by the urea denaturation study. CONCLUSIONS: **Arginine** perturbs the tertiary and quaternary structure of **alpha-crystallin** and enhances the dynamics of the subunit assembly leading to enhanced chaperone-like activity. Thus, in addition to size, surface hydrophobicity, and charge distribution, the dynamics of the subunit assembly appears to be one of the critical factors that can modulate the chaperone activity.

Descriptors: \***Arginine** --pharmacology--PD; \*Molecular Chaperones --metabolism--ME; \***alpha-Crystallins**--drug effects--DE; Animals; Cattle; Disulfides; Dithiothreitol; Fluorescent Dyes; Lens, Crystalline--chemistry --CH; Protein Subunits--chemistry--CH; Protein Subunits--metabolism--ME; Recombinant Proteins--chemistry--CH; Recombinant Proteins--drug effects --DE; Recombinant Proteins--metabolism--ME; Solubility; Spectrometry, Fluorescence; **alpha-Crystallins**--chemistry--CH; **alpha-Crystallins** --metabolism--ME

CAS Registry No.: 0 (Disulfides); 0 (Fluorescent Dyes); 0 (Molecular Chaperones); 0 (Protein Subunits); 0 (Recombinant Proteins); 0 (**alpha-Crystallins**); 3483-12-3 (Dithiothreitol); 74-79-3 (**Arginine**)

Record Date Created: 20050413

Record Date Completed: 20060413

Date of Electronic Publication: 20050401

**Arginine hydrochloride enhances the dynamics of subunit assembly and the chaperone-like activity of alpha-crystallin.**

PURPOSE: **Alpha-crystallin**, a major eye lens protein, bears homology with small heat shock proteins (sHsps) and exhibits...

...have earlier demonstrated similar enhancement of chaperone-like activity using biologically compatible solutes such as **arginine hydrochloride** and aminoguanidine. The purpose of the present study is to get an insight into the mechanism of the **arginine** induced enhancement of chaperone-like

activity of alpha- **crystallin** . METHODS: The effect of **arginine hydrochloride** on the chaperone-like activity of alpha- **crystallin** at 25 degrees C was studied using DTT induced aggregation of insulin as a model system. Changes in the accessibility of the thiol group near the end of the alpha- **crystallin** domain in the absence and the presence of **arginine hydrochloride** were studied using dithiobisnitrobenzoic acid. Fluorescence resonance energy transfer studies were performed to investigate changes in the dynamics of the subunit assembly. Urea induced denaturation studies of alpha- **crystallin** were carried out to investigate structural destabilization of alpha- **crystallin** , if any, in the presence of **arginine hydrochloride** . RESULTS: **Arginine hydrochloride** increases the chaperone-like activity of alpha- **crystallin** several fold towards DTT induced aggregation of insulin at room temperature. Our study shows that...

... and the rate of accessibility of the thiol group are increased in the presence of **arginine** . Fluorescence resonance energy transfer experiments show that **arginine hydrochloride** significantly increases the subunit exchange between the oligomers of alpha- **crystallin** . **Arginine** induced structural perturbation and loosening of subunit assembly of alpha- **crystallin** leads to overall destabilization of the protein as reflected by the urea denaturation study. CONCLUSIONS: **Arginine** perturbs the tertiary and quaternary structure of alpha- **crystallin** and enhances the dynamics of the subunit assembly leading to enhanced chaperone-like activity. Thus...

Descriptors: \***Arginine** --pharmacology--PD; \*Molecular Chaperones --metabolism--ME; \*alpha-Crystallins--drug effects--DE .

Chemical Name: Disulfides; Fluorescent Dyes; Molecular Chaperones; Protein Subunits; Recombinant Proteins; alpha-Crystallins; Dithiothreitol; **Arginine**

9/9,K/20 (Item 3 from file: 155)

DIALOG(R) File 155: MEDLINE(R)

(c) format only 2006 Dialog. All rts. reserv.

14316050 PMID: 12761397

**Structural perturbation and enhancement of the chaperone-like activity of alpha- crystallin by arginine hydrochloride .**

Srinivas Volety; Raman Bakthisaran; Rao Kunchala Sridhar; Ramakrishna Tangirala; Rao Ch Mohan

Centre for Cellular & Molecular Biology, Hyderabad 500 007, India.

Protein science - a publication of the Protein Society (United States)

Jun 2003, 12 (6) p1262-70, ISSN 0961-8368--Print Journal Code: 9211750

Publishing Model Print

Document type: Journal Article

Languages: ENGLISH

Main Citation Owner: NLM

Record type: MEDLINE; Completed

Subfile: INDEX MEDICUS

Structural perturbation of alpha- **crystallin** is shown to enhance its molecular chaperone-like activity in preventing aggregation of target proteins. We demonstrate that **arginine** , a biologically compatible molecule that is known to bind to the peptide backbone and negatively charged side-chains, increases the chaperone-like activity of calf eye lens alpha- **crystallin** as well as recombinant human alphaA- and alphaB-crystallins. **Arginine** -induced increase in the chaperone activity



is more pronounced for alphaB- crystallin than for alphaA- crystallin . Other guanidinium compounds such as aminoguanidine hydrochloride and guanidine hydrochloride also show a similar effect, but to different extents. A point mutation, R120G, in alphaB- crystallin that is associated with desmin-related myopathy, results in a significant loss of chaperone-like activity. Arginine restores the activity of mutant protein to a considerable extent. We have investigated the effect of arginine on the structural changes of alpha- crystallin by circular dichroism, fluorescence, and glycerol gradient sedimentation. Far-UV CD spectra show no significant changes in secondary structure, whereas near-UV CD spectra show subtle changes in the presence of arginine . Glycerol gradient sedimentation shows a significant decrease in the size of alpha- crystallin oligomer in the presence of arginine . Increased exposure of hydrophobic surfaces of alpha- crystallin , as monitored by pyrene-solubilization and ANS-fluorescence, is observed in the presence of arginine . These results show that arginine brings about subtle changes in the tertiary structure and significant changes in the quaternary structure of alpha- crystallin and enhances its chaperone-like activity significantly. This study should prove useful in designing strategies to improve chaperone function for therapeutic applications.

Descriptors: \*Arginine --pharmacology--PD; \*Crystallins--chemistry--CH; Animals; Cattle; Centrifugation, Density Gradient; Circular Dichroism; Crystallins--metabolism--ME; Dithiothreitol; Guanidine--pharmacology--PD; Insulin--chemistry--CH; Insulin--metabolism--ME; Protein Conformation --drug effects--DE; Pyrenes--chemistry--CH; Solubility; Spectrometry, Fluorescence; Time Factors

CAS Registry No.: 0 (Crystallins); 0 (Pyrenes); 11061-68-0 (Insulin); 113-00-8 (Guanidine); 129-00-0 (pyrene); 3483-12-3 (Dithiothreitol); 74-79-3 (Arginine)

Record Date Created: 20030522

Record Date Completed: 20041005

#### **Structural perturbation and enhancement of the chaperone-like activity of alpha- crystallin by arginine hydrochloride .**

Structural perturbation of alpha- crystallin is shown to enhance its molecular chaperone-like activity in preventing aggregation of target proteins. We demonstrate that arginine , a biologically compatible molecule that is known to bind to the peptide backbone and negatively charged side-chains, increases the chaperone-like activity of calf eye lens alpha- crystallin as well as recombinant human alphaA- and alphaB-crystallins. Arginine -induced increase in the chaperone activity is more pronounced for alphaB- crystallin than for alphaA- crystallin . Other guanidinium compounds such as aminoguanidine hydrochloride and guanidine hydrochloride also show a similar effect, but to different extents. A point mutation, R120G, in alphaB- crystallin that is associated with desmin-related myopathy, results in a significant loss of chaperone-like activity. Arginine restores the activity of mutant protein to a considerable extent. We have investigated the effect of arginine on the structural changes of alpha- crystallin by circular dichroism, fluorescence, and glycerol gradient sedimentation. Far-UV CD spectra show no significant...

... in secondary structure, whereas near-UV CD spectra show subtle changes in the presence of arginine . Glycerol gradient sedimentation shows a significant decrease in the size of alpha- crystallin oligomer in the presence of arginine . Increased exposure of hydrophobic surfaces of alpha- crystallin , as monitored by pyrene-solubilization and